GEOMETRICAL
CONIC SECTIONS

ar V
19530

39 - B

Cornell University Library
arV19530

A geometrical treatise on conic sections

3 1924 031 271 509
olin,anx

Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.

http://www.archive.org/details/cu31924031271509

A GEOMETRICAL TREATISE

ON

CONIC SECTIONS.



A

GEOMETRICAL TREATISE

ON

CONIC SECTIONS.

WITH NUMEROUS EXAMPLES.

For the Use of Schools and Students in the Universities.

By the REV. W. H. DREW, M.A.
ST. JOHN'S COLLEGE, CAMBRIDGE,
PROFESSOR OF MATHEMATICS IN KING'S COLLEGE, LONDON.

FOURTH EDITION.

London and Cambridge:
MACMILLAN AND CO.
MDCCCLXIX.

[*The Right of Translation and Reproduction is reserved.*]

~~8417 A 30.~~

CORNELL
UNIVERSITY
LIBRARY

Kww

LONDON:
R. CLAY, SONS, AND TAYLOR, PRINTERS,
BREAD STREET HILL.

CONIC SECTIONS.

INTRODUCTION.

1. **Def.** The curve traced out by a point, which moves in such a manner that its distance from a given fixed point continually bears the same ratio to its distance from a given fixed line, is called a *Conic Section.*

The fixed point is called the *Focus,* and the fixed line the *Directrix.*

Thus if S be the focus, and KK' the directrix, and P a point from which PM is drawn at right angles to the directrix, the curve traced out by P will be a *Conic Section,* provided P move in such manner that SP always bears the same ratio to PM.



(1.) When the distance from the fixed point is equal to the distance from the fixed line, that is, when SP is equal to PM, the *Conic Section* is called a *Parabola.*

(2.) When the distance from the fixed point is less than the distance from the fixed line, that is, when the ratio which

SP bears to *PM* is less than unity, the *Conic Section* is called an *Ellipse.*

(3.) When the distance from the fixed point is greater than the distance from the fixed line, that is, when the ratio which *SP* bears to *PM* is greater than unity, the *Conic Section* is called an *Hyberbola.*

2. The reason of the term *Conic Sections* being applied to these curves is that, when a *Cone* is intersected by a plane surface, the boundary of the section so formed will, *in general,* be one or other of these curves.

I purpose to investigate the properties of the *Conic Sections* from the definitions given above, and afterwards to show in what manner a *Cone* must be divided by a plane in order that the curve of intersection may be a *Parabola, Ellipse,* or *Hyperbola.*

CHAPTER I.

THE PARABOLA.

PROP. I.

3. The focus and directrix of a parabola being given, to find any number of points on the curve.



Let S be the focus, and KK' the directrix.

Draw XSx at right angles to the directrix, and bisect the line SX in A; then

$$\text{since } AS = AX,$$

$$\therefore A \text{ is a point on the curve.}$$

The point A is called the *Vertex*, and the line Ax, with respect to which the curve is evidently symmetrical, is called the *Axis*.

On the directrix take any point M; join SM; and draw MP at right angles to the directrix.

At the focus S make the angle MSP equal to the angle SMP; then

$$SP = PM,$$

$$\therefore P \text{ is a point on the curve.}$$

So by taking any number of points, M', M'', on the directrix, we may obtain as many points, P', P'', on the curve as we please, and the line which passes through A and all these points will be the parabola whose focus is S and directrix KK'.

Cor. 1. As M is taken further away from the point X, the line SM and the angles SMP, MSP, and, consequently, the lines SP and PM, continually increase. Hence, since XM and MP increase together, the curve recedes at the same time both from the axis and directrix; and since the angle SMP can never exceed a right angle, and the lines SP and MP will therefore always meet, it is evident that there is no limit to the distance to which the curve may extend on both sides of the axis.

Cor. 2. The parabola may be described practically in the following manner.



Let S be the focus and KX be the directrix; and let a rigid bar QM, having a string of the same length as itself fastened at one end Q, be made to slide parallel to the axis with the other end M on the directrix; then if the other end of the string be fastened at the focus, and the string be kept stretched by means of the point of a pencil at P, in contact with the bar, since SP will always be equal to PM, it is evident that the point P will trace out the parabola.

PROP. II.

4. The distance of any point inside the parabola from the focus is less than its distance from the directrix; and the distance of any point outside the parabola from the focus is greater than its distance from the directrix.



(1.) Let Q be a point inside the parobola.

Draw QM at right angles to the directrix, meeting the parobola in P; join SP; then

$$\text{since } SP = PM,$$
$$\therefore SP \text{ and } PQ = QM.$$
$$\text{But } SP \text{ and } PQ > SQ,$$
$$\therefore QM > SQ.$$

(2.) Let Q be a point outside the parabola.

Draw MQ at right angles to the directrix, and produce it to meet the parabola in P; join SP; then

$$\text{since } SQ \text{ and } QP > SP,$$
$$\text{and } SP = PM,$$
$$\therefore SQ \text{ and } QP > PM,$$
$$\therefore SQ > QM.$$

COR. Conversely a point will be inside or outside the parabola according as its distance from the focus is less or greater than its distance from the directrix.

5. DEF. The line PN (*see fig. Prop.* III.) drawn at right angles to the axis from the point P in the curve is called the *Ordinate* of the point P, and the line AN the *Abscissa.* The double ordinate BC drawn through the focus, and terminated both ways by the curve, is called the *Latus Rectum.*

Prop. III.

The Latus Rectum $BC = 4AS$.



$PN^2 + SN^2 = PS^2$
$= XN^2$
$\therefore y^2 + (x-a)^2 = (x+a)^2$
$y^2 = 4ax$

Draw BK at right angles to the directrix.

$$\text{Then } SB = BK = SX = 2AS,$$
$$\therefore BC = 4AS.$$

6. Def. If a point P' be taken on the parabola (*see fig. Prop.* IV.) near to P, and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P, is called the *Tangent* to the parabola at the point P.

Prop. IV.

If the tangent to the parabola at any point P intersect the directrix in the point Z; then SZ will be at right angles to SP.

Let P' be a point on the parabola near to P.



Draw the chord PP', and produce it to meet the directrix in Z; join SZ.

Draw PM, $P'M'$ at right angles to the directrix; join SP, SP'; and produce PS to meet the parabola in Q.

Then, since the triangles ZMP, $ZM'P'$ are similar,

$$\therefore ZP : ZP' :: MP : M'P',$$
$$:: SP : SP',$$

$\therefore$ SZ bisects the angle $P'SQ$. (*Euclid*, VI. *Prop.* A.)

Now when P' is indefinitely near to P, and PP' becomes the tangent at the point P, the angle PSP' becomes indefinitely small, while the angle QSP' approaches two right angles, and therefore the angle $P'SZ$, which is half of the angle $P'SQ$, becomes ultimately a right angle.

Hence, when PZ is the tangent,

the angle ZSP is a right angle,
or SZ is perpendicular to SP.

Cor. Conversely, if SZ be drawn at right angles to SP, meeting the directrix in Z, and PZ be joined, PZ will be a tangent at P.

PROP. V.

7. The tangent at any point P of a parabola bisects the angle between the focal distance SP, and the perpendicular PM on the directrix.



Let the tangent at P meet the directrix in the point Z; join SZ; then since the angle ZSP is a right angle, (*Prop.* IV.)

$$\cdot\ ZS^2 + SP^2 = PZ^2.$$

$$\text{Also } ZM^2 + MP^2 = PZ^2.$$

$$\therefore ZS^2 + SP^2 = ZM^2 + MP^2.$$

$$\text{But } SP = PM,$$

$$\therefore ZS = ZM.$$

Now in the triangles ZPS, ZPM,

$\because ZP, PS = ZP, PM$, each to each,

and $ZS = ZM$,

. the angle SPZ = the angle MPZ;

or PZ bisects the angle SPM.

COR. 1. If ZP be produced to R, then the angle SPR = the angle MPR.

COR. 2. It is evident that the tangent at the vertex A is perpendicular to the axis.

PROP. VI.

8. The tangents at the extremities of a focal chord intersect at right angles in the directrix.

Let PSQ be a focal chord, and let the tangent at P meet the directrix in Z.

Join SZ; then

the angle ZSP is a right angle, (*Prop.* IV.)

and ∴ also the angle ZSQ is a right angle,

∴ ZQ is the tangent at Q, (*Prop.* IV. *Cor.*)

or the tangents at the extremities of the focal chord PSQ intersect in the directrix.

Again, draw PM, QM' at right angles to the directrix; then

since MP, PZ = SP, PZ, each to each,

and the angle MPZ = the angle SPZ,

∴ the angle MZP = the angle SZP,

∴ the angle SZP is half of the angle SZM.

So the angle SZQ is half of the angle SZM',

∴ the angle PZQ is half of the two SZM and SZM.

But the angles SZM and SZM' = two right angles,

∴ the angle PZQ is a right angle,

or the tangents at the extremities of a focal chord intersect *at right angles* in the directrix.

Prop. VII.

9. If the tangent at any point P of a parabola meet the axis produced in the point T, and PN be the ordinate of the point P, then $NT = 2AN$.



Join SP, and draw PM at right angles to the directrix; then

$$\because \text{ the angle } SPT = \text{ the angle } MPT = \text{ the angle } STP,$$

$$\therefore\ ST = SP.$$

$$\text{But } SP = PM = XN,$$

$$\therefore\ ST = XN.$$

$$\text{But } AS = AX,$$

$$\therefore \text{ the remainder } AT = \text{ the remainder } AN,$$

$$\therefore\ NT = 2AN.$$

Def. The line NT is called the *Subtangent*.

10. Def. The line PG, drawn at right angles to PT, is called the *Normal* at the point P, and NG the *Subnormal*.

Prop. VIII.

If the normal at the point P of a parabola meet the axis in the point G, then $NG = 2AS$.

Since the angle SPG = the complement of the angle SPT,
and the angle SGP = the complement of the angle STP,
and also the angle SPT = the angle STP, (*Prop.* VII.)
$\therefore$ the angle SPG = the angle SGP,
$\therefore SG = SP$.
But $SP = PM = XN$,
$\therefore SG = XN$.
Taking away the common part SN,
the remainder $NG = SX = 2AS$.

Prop. IX.

11. If PN be an ordinate to the parabola at the point P; then $PN^2 = 4AS \,.\, AN$.

Since TPG is a right angle, and PN perpendicular to TG;
$\therefore PN$ is a mean proportional between TN and NG;
or $PN^2 = TN \,.\, NG$. (*Euclid*, VI. 8 *Cor.*)
But $TN = 2AN$, (*Prop.* VII.)
and $NG = 2AS$, (*Prop.* VIII.)
$\therefore PN^2 = 4\,AS \,.\, AN$.

Prop. X.

12. If the tangent at any point P intersect the tangent at the vertex in Y, then SY will bisect PT at right angles, and will be a mean proportional between SA and SP.

Draw PN at right angles to the axis; then
since AY is parallel to PN,
$\therefore TY : YP :: TA : AN$.
But $AT = AN$, (*Prop.* VII.)
$\therefore TY = PY$;
and $\because SY, YP = SY, YT$, each to each,
and $SP = ST$, (*Prop.* VII.)
$\therefore$ the angle SYP = the angle SYT,
$\therefore SY$ is perpendicular to PT.

Again, since TYS is a right angle, and YA perpendicular to ST,

∴ SY is a mean proportional between ST and SA;

or $SY^2 = ST \,.\, SA$. (*Euclid*, VI. 8 *Cor.*)

But $ST = SP$, (*Prop.* VII.)

$\therefore\ SY^2 = SP \,.\, SA$.

Cor. If PM be drawn at right angles to the directrix, and MY be joined, then

since SP, $PY = MP$, PY, each to each,

and the angle $SPY =$ the angle MPY, (*Prop.* V.)

∴ the angle $SYP =$ the angle MYP,

∴ SY and YM are in the same straight line.

Prop. XI.

13. To draw a pair of tangents to a parabola from an external point.



Let O be the given external point.

Join OS, and with centre O and radius OS describe a circle, cutting the directrix in M and M', which it will always do, on whichever side of the directrix O is situated, since O is nearer to the directrix than to the focus. (*Prop.* II.)

Draw MQ and $M'Q'$ parallel to the axis meeting the parabola in Q and Q'.

Join OQ, OQ'; these will be the tangents required.

Join SQ and SQ'; then

$\because$ OQ, $QS = OQ$, QM, each to each,

and $OS = OM$,

$\therefore$ the angle $OQS =$ the angle OQM,

$\therefore$ OQ is the tangent at Q. (*Prop.* V.)

So OQ' is the tangent at Q'.

PROP. XII.

14. If from a point O a pair of tangents OQ and OQ' be drawn to a parabola, the triangles OSQ, OSQ' will be similar, and OS will be a mean proportional between SQ and SQ'.

Join SM, cutting OQ at right angles (*Prop.* X. *Cor.*) in the point Y; then

since the angle $SQO =$ the angle MQO, (*Prop.* V.)

and the angle $MQO =$ the angle SMM',

each of these angles being the complement of the angle QMY,

$\therefore$ the angle $SQO =$ the angle SMM'.

But the angle SMM' at the circumference is half the angle SOM' at the centre, and is therefore equal to the angle SOQ'.

$\therefore$ the angle $SQO =$ the angle SOQ'.

So the angle $SOQ =$ the angle $SQ'O$,

$\therefore$ the remaining angle $OSQ =$ the remaining angle OSQ'.

And therefore the triangle OSQ is similar to the triangle OSQ'.

$$\therefore SQ : SO :: SO : SQ',$$

$$\therefore SQ \, . \, SQ' = SO^2.$$

or SO is a mean proportional between SQ and SQ'.

Prop. XIII.

15. If a pair of tangents OQ, OQ' be drawn to a parabola, and OV be drawn parallel to the axis meeting QQ' in V, then QQ' shall be bisected in V.



Draw QM, $Q'M'$ at right angles to the directrix.

Join OM, OM'; and let OV meet MM' in Z.

Then, since $OM = OM'$, (*Prop.* XI.)

$\therefore$ the angle $OMZ =$ the angle $OM'Z$,

and the angle $OZM =$ the angle OZM',

and the side OZ is common to the triangles OZM, OZM',

$$\therefore MZ = M'Z.$$

And because the lines QM, ZV, $Q'M'$ are parallel,

$$\therefore QV : Q'V :: MZ : M'Z.$$

$$\text{But } MZ = M'Z,$$

$$\therefore QV = Q'V,$$

$$\therefore QQ' \text{ is bisected in } V.$$

Prop. XIV.

16. If from a point O a pair of tangents OQ, OQ', be drawn to a parabola, and OV be drawn parallel to the axis meeting the parabola in P, and QQ' in V, then the tangent at P will be parallel to QQ' and OV will be bisected in P.



Draw the tangent RPR' meeting OQ, OQ' in R and R'.

Join PQ, and draw RW parallel to the axis, meeting PQ in W;

Then, by the last Proposition,

$$PW = WQ.$$

And because RW is parallel to OP,

$$\therefore OR : RQ :: PW : WQ.$$

$$\text{But } PW = WQ,$$

$$\therefore OR = RQ;$$

$$\text{so } OR' = R'Q',$$

$$\therefore OR : RQ :: OR' : R'Q',$$

$$\therefore RR' \text{ is parallel to } QQ'.$$

Again, since PR is parallel to QV,

$$\therefore OP : PV :: OR : RQ.$$

$$\text{But } OR = RQ,$$

$$\therefore OP = PV.$$

Cor. From this it is manifest that if any number of parallel chords be drawn in a parabola, their middle points will all lie on the line parallel to the axis which passes through the point where the tangent drawn parallel to the chords meets the parabola.

Def. Any line PV, drawn from a point P in the parabola parallel to the axis, is called a *Diameter*.

The point P is called the *Vertex* of the diameter PV; and the tangent at P the *Tangent at the Vertex.*

The diameter consequently bisects all chords parallel to the tangent at the vertex, and the tangents at the extremities of any chord will intersect in the diameter corresponding to that chord.

Def. A line QV, drawn parallel to the tangent at P from a point Q in the curve, is called the *Ordinate* to the diameter PV.

Prop. XV.

17. If QV be an ordinate to the diameter PV, then $QV^2 = 4 . SP . PV.$



Produce QV to meet the parabola in Q'; and draw the tangents QO, $Q'O$, meeting VP produced in the point O. (*Prop.* XIV.)

Also let the tangent at P meet OQ in R, and join SP, SR, and SQ. Now since from the point R two tangents RP, RQ are drawn to the parabola, the triangle RPS is similar to the triangle RSQ, (*Prop.* XII.)

$\therefore$ the angle SRP = the angle SQR.

But the angle SQR = the angle STQ, (*Prop.* VII.)

= the angle POR,

$\therefore$ the angle SRP = the angle POR,

and the angle SPR = the angle OPR, (*Prop.* V. *Cor.* 1.)

$\therefore$ the remaining angle RSP = the remaining angle ORP,

$\therefore$ the triangle SPR is similar to the triangle POR,

$$\therefore SP : PR :: PR : PO,$$

$$\therefore PR^2 = SP \,.\, PO,$$

$$= SP \,.\, PV. \quad (\textit{Prop.}\ \text{XIV.})$$

Again, since QV is parallel to PR,

$$\therefore QV : PR :: OV : OP.$$

But $OV = 2\,OP$, (*Prop.* XIV.)

$$\begin{aligned} \therefore\ QV &= 2\,PR, \\ \therefore\ QV^2 &= 4\,PR^2, \\ &= 4\,SP\,.\,PV. \end{aligned}$$

18. Def. The double ordinate to the diameter PV, drawn parallel to the tangent at P, and passing through the focus, is called the *Parameter* of *the diameter PV.*

Prop. XVI.

The parameter of the diameter $\overset{QQ'}{\cancel{PV}} = 4\,.\,SP.$



Draw QSQ' through the focus parallel to the tangent at P, and let the tangent at P meet the axis produced in T; then

$$QV^2 = 4\,SP\,.\,PV. \quad (\textit{Prop.}\ \text{XV.})$$

$$\text{But } PV = ST = SP, \quad (\textit{Prop.}\ \text{VII.})$$

$$\begin{aligned} \therefore\ QV^2 &= 4\,SP^2; \\ \text{or } QV &= 2\,SP, \\ \therefore\ QQ' &= 4\,SP. \end{aligned}$$

PROP. XVII.

19. If two chords of a parabola intersect one another, the rectangles contained by their segments are in the ratio of the parameters of the diameters which bisect the chords.



Let the chords Qq, $Q'q'$ intersect one another in the point O.

Bisect Qq, $Q'q'$ in V and V'; and draw the diameters PV, $P'V'$ parallel to the axis.

Also, through O draw OR parallel to PV; and through R draw RW parallel to QV.

Now, since Qq is divided equally in V and unequally in O,

$$\therefore QO \cdot Oq = QV^2 - OV^2, \quad (\textit{Euclid}, \text{ II. } 5)$$

$$= QV^2 - RW^2,$$

$$= 4SP \cdot PV - 4SP \cdot PW, \quad (\textit{Prop.} \text{ XV.})$$

$$= 4SP \cdot RO.$$

So $Q'O \cdot Oq' = 4SP' \cdot RO$.

Hence $QO \cdot Oq : Q'O \cdot Oq' :: 4SP : 4SP'$.

By *Euclid*, II. 6, the same may be proved to be true if the point O be without the parabola.

PROP. XVIII.

20. If from an external point O a pair of tangents OQ, OQ' be drawn to the parabola, and the chord QQ' be joined, the area of the figure bounded by QQ' and the curve is two-thirds of the triangle QOQ'.



Draw the diameter OV meeting the curve in P; and let the tangent at P meet OQ, OQ' in R and R'.

Join QP, $Q'P$; then

$$\text{since } OR = RQ,$$

$$\therefore \text{ the triangle } OPR = \tfrac{1}{2} \text{ the triangle } OPQ,$$

$$= \tfrac{1}{2} \text{ the triangle } VPQ.$$

So the triangle $OPR' = \frac{1}{2}$ the triangle VPQ',

$\therefore$ the triangle $ORR' = \frac{1}{2}$ the triangle PQQ'.

Again, if through R and R' we draw the diameters Rp, $R'p'$; and at the points p and p' draw the tangents $rpr_{,}$, $r'p'r'_{,}$, we can prove in the same manner as before that

the triangle $Rrr_{,} = \frac{1}{2}$ the triangle QpP,

and the triangle $R'r'r'_{,} = \frac{1}{2}$ the triangle $Q'p'P$.

Continuing in this manner to form new triangles by drawing diameters at the points r, $r_{,}$, and r', $r'_{,}$, and tangents at the points where these diameters meet the curve, we can prove that the exterior triangles formed by the tangents are the halves of the interior triangles formed by joining the points of contact with the extremities of the chords.

And the same will hold however the number of the triangles be increased.

Hence the sum of all the exterior triangles will be equal to half the sum of all the interior angles.

Now when the number of the triangles is increased indefinitely, the sum of the exterior triangles will represent the exterior figure $OQPQ'$, and the sum of the interior triangles the area of the interior figure QPQ. Hence

the area of the figure $OQPQ' = \frac{1}{2}$ the area of the figure QPQ'

$\therefore$ area of the figure $OQPQ' = \frac{1}{3}$ the area of triangle QOQ',

$\therefore$ area of the figure $QPQ' = \frac{2}{3}$ the area of triangle QOQ'.

21. Def. If with a point O on the normal at P as centre and OP as radius, a circle be described touching the parabola at P and cutting it in Q; then when the point Q is made to approach indefinitely near to P, the circle is called the *Circle of Curvature* at the point P. (*See fig. Prop.* XIX.)

Prop. XIX.

The chord of the circle of curvature, at a point P of a parabola, drawn parallel to the axis $= 4SP$.

Let PT be the tangent, and PG the normal at the point P.

With centre O and radius OP describe a circle cutting the parabola in the point Q.

Draw RQX parallel to the axis meeting the circle in X and the tangent at P in R.

Also draw QV parallel to PR, and PW parallel to the axis; then

since RP touches the circle at P,

$$\therefore RQ \cdot RX = PR^2. \quad (\textit{Euclid}, \text{ III. } 36.)$$



$$\text{But } PR^2 = QV^2 = 4SP \,.\, PV, \quad (\textit{Prop.}\ \text{XV.})$$
$$\therefore RQ \,.\, RX = 4SP \,.\, PV.$$
$$\text{But } RQ = PV.$$
$$\therefore RX = 4SP.$$

Now when the circle becomes the circle of curvature at P, the points R and Q move up to and coincide with P, and the lines RX and PW become equal.

Hence the chord of the circle of curvature parallel to the axis = $4SP$.

Cor. 1. If PU be the diameter of the circle of curvature, and PF the chord through the focus; then

$$\text{since the angle } FPU = \text{the angle } WPU, \quad (\textit{Prop.}\ \text{VIII.})$$
$$\therefore PF = PW = 4SP.$$

Cor. 2. If SY be drawn at right angles to PT; then

$$\text{the triangle } PFU \text{ is similar to } SYP,$$
$$\therefore PU : PF :: SP : SY,$$
$$\text{or } PU : 4SP :: SP : SY.$$

Prop. XX.

If QVQ' be any ordinate to the diameter PV, the circle described through the three points P, Q, Q' will intersect the parabola in a fourth point, which depends only upon the position of P.

Draw the ordinate PN, and produce it to meet the parabola in P'; then,

$$\text{since the subtangent} = 2 \,.\, AN. \quad (\textit{Prop.}\ \text{VII.})$$

The tangents at P and P' will meet the axis in the same point T.

Draw PR parallel to TP', meeting the parabola in R, and QQ' in O; then

$$PO \,.\, OR : QO \,.\, OQ' :: SP' : SP. \quad (\textit{Prop.}\ \text{XVII.})$$



$$\text{But } SP = ST = SP', \text{ (\textit{Prop.} VII.)}$$

$$\therefore PO \,.\, OR = QO \,.\, OQ'.$$

Hence by the converse of *Euclid* III. *Prop.* 22, the point R is on the circle which passes through P, Q, Q'.

Cor. 1. Since TP and TP' are equally inclined to the axis, the lines QQ', PR, which are parallel respectively to TP and TP', are also equally inclined to the axis.

Cor. 2. When the point V is brought indefinitely near to P, QQ' coincides with the tangent to the parabola at P, and becomes also a tangent to the circle at P, since Q and Q' are indefinitely near to each other. The circle therefore becomes the circle of curvature at the point P.

Hence if PR be drawn parallel to the tangent at P', or be equally inclined to the axis with PT, it will meet the parabola in the point where the circle of curvature at P intersects the parabola.

PROBLEMS ON THE PARABOLA.

1. The diameter of the circle described about the triangle BAC is equal to $5AS$. (*See fig. Prop.* III.)

2. If from the point G, GK be drawn at right angles to SP, then $PK = 2AS$. (*See fig. Prop.* VII.)

3. If the triangle SPG is equilateral, then SP is equal to the latus rectum. (*See fig. Prop.* VII.)

4. PQ is a common tangent to a parabola and the circle described on the latus rectum as diameter; prove that SP and SQ make equal angles with the latus rectum.

5. Prove that $PY . PZ = SP^2$, and that $PY . YZ = AS . SP$. (*See fig. Prop.* VII.)

6. If PL be drawn at right angles to AP, meeting the axis in L, and PN be the ordinate of P, then $NL = 4AS$.

7. The tangent at any point P of a parabola meets the directrix and latus rectum produced in points equally distant from the focus.

8. Prove that $NY = TY$, and that $TP . TY = TS . TN$. (*See fig. Prop.* VII.)

9. If a circle be described about the triangle SPN, the tangent to it from $A = \frac{1}{2} PN$. (*See fig. Prop.* VII.)

10. If the ordinate of a point P bisect the subnormal of P', the ordinate of P is equal to the normal of P'.

11. If from any point on the tangent to a parabola a line be drawn touching the parabola, the angle between this line and the line to the focus from the same point is constant.

12. A circle and parabola have the same vertex and axis. $BA'C$ is the double ordinate of the parabola which touches the circle at A', the extremity of the diameter through the vertex A. PP' is any other ordinate of the parabola parallel to this, meeting the axis in N, and AB produced in R; prove that the rectangle $RP \,.\, RP'$ is proportional to the square of the tangent drawn from N to the circle.

13. Draw a parabola to touch a given circle at a given point, and such that its axis may touch the same circle in another given point.

14. If from the point of contact of a tangent to a parabola a chord be drawn, and another line be drawn parallel to the axis meeting the chord, tangent, and curve, this line will be divided by them in the same ratio as it divides the chord.

15. If the diameter PV meet the directrix in O, and the chord drawn through the focus parallel to the tangent at P in V, prove that $VP = PO$.

16. Prove that the locus of the intersection of a diameter PV with the chord drawn through the focus parallel to the tangent at P is a parabola.

17. If a circle and parabola have a common tangent at P, and intersect in Q and R; and QV, UR be drawn parallel to the axis of the parabola meeting the circle in V and U respectively, then VU is parallel to the tangent at P.

18. AB and AC are two lines at right angles to each other. From a fixed point C on AC, CR is drawn parallel to AB. On AR, produced if necessary, P is taken such that the perpendicular PN upon AB is equal to CR. Prove that the curve traced out by P is a parabola.

19. If from a point P of a circle PC be drawn to the centre; and R be the middle point of the chord PQ drawn parallel to a fixed diameter ACB, then the curve traced out by the intersection of CP and AR is a parabola.

20. If two equal tangents OQ, OQ', be cut by a third tangent, their alternate segments are equal.

21. E is the centre of the circle described about the triangle OQQ'. Prove that the circle described about the triangle QEQ' will pass through the focus. (*See fig. Prop.* XIII.)

22. PSp is any focal chord of a parabola. Prove that AP, Ap will meet the latus rectum in two points Q, q, whose distances from the focus are equal to the ordinates of p and P.

23. PSp is a focal chord of a parabola, RDr the directrix meeting the axis in D; and Q any point on the curve. Prove that if QP, Qp be produced to meet the directrix in R, r, half the latus rectum is a mean proportional between DR, Dr.

24. OP and OQ are two tangents to a parabola. On QO produced, OQ' is taken equal to OQ; prove that $OS . PQ' = OP . OQ$.

25. If QD be drawn at right angles to the diameter PV, then $QD^2 = 4AS . PV$.

26. If through any point O on the axis of a parabola a chord POQ be drawn, and PM, QN be the ordinates of the points P and Q, prove that $AM . AN = AO^2$.

27. If AP and AQ be drawn at right angles to each other from the vertex of a parabola, and PM, QN be the ordinates of P and Q, prove that the latus rectum is a mean proportional between AM and AN.

28. OAP is the sector of a circle whose centre is O. If the radius OA remain fixed while the angle AOP changes the centre of the circle inscribed in the sector, AOP will trace out a parabola.

29. QSQ' is a focal chord parallel to AP; PN, QM, $Q'M'$ are the ordinates of P, Q, and Q'. Prove that $SM^2 = AM . AN$ and that $MM' = AP$.

30. PQ, PQ' are drawn from any point P cutting the ordinates $Q'V'$, QV in R' and R, prove that VR is to $V'R'$ in the triplicate ratio of QV to $Q'V'$.

31. On a chord of a parabola as diameter a circle is described cutting the parabola again in two points. If these points be joined, the portion of the axis between the two chords is equal to the latus rectum.

32. If OQ, OQ' be a pair of tangents to a parabola, and the chord QQ' be a normal to the curve at Q, then OQ is bisected by the directrix.

33. Two equal parabolas having the same focus and their axes in contrary directions intersect at right angles.

34. The radius of curvature at the extremity of the latus rectum is equal to twice the normal.

35. If from any point P of a parabola PF and PH be drawn making equal angles with the normal PG, then $SG^2 = SF \cdot SH$.

36. If a triangle be inscribed in a parabola, the points when the sides produced meet the tangents at the opposite angles are in the same straight line.

37. If the tangents OQ, OQ' be cut by a third tangent in R, R', prove that

$$OR : RQ :: R'Q' : OR'.$$

38. If from the vertex of a parabola chords be drawn at right angles to one another, and on them a rectangle be described, the curve traced out by the further angle is a parabola.

39. Prove that $2PY$ is a mean proportional between AP and the chord of the circle of curvature at the point P of the parabola drawn through the vertex A. (*See fig. Prop.* VII.)

40. If a circle described upon the chord of a parabola as diameter meet the directrix it also touches it, and all chords for which this is possible intersect in a point.

41. If a parabola *roll* upon another equal parabola, the vertices originally coinciding, the focus traces out the directrix.

42. The circle of curvature at the extremity of the latus rectum intersects the parabola on the diameter of curvature passing through the point of contact.

CHAPTER II.

THE ELLIPSE.

22. Def. The *Ellipse* is the curve traced out by a point which moves in such a manner that its distance from a given fixed point continually bears the same ratio, *less than unity,* to its distance from a given fixed line. (*See Introduction.*)

Prop. I.

The focus and directrix of an ellipse being given, to find any number of points on the curve.

Let S be the focus, and MX the directrix.

Draw SX at right angles to the directrix, and divide SX in the point A, so that SA may be to AX in the given fixed ratio less than unity; then

A is a point on the curve.

On XS produced take a point A', such that

$$SA' : A'X :: SA : AX;$$

then A' will also be a point on the curve.

On the directrix take *any* point M; and through M and S draw the line $MYSY'$ meeting AY and $A'Y'$, drawn at right angles to AA' in the points Y and Y'.

On YY' as diameter describe a circle, and draw MPP' parallel to AA', cutting the circle in the points P and P';

P and P' will be points on the ellipse.



Join PY, PY', SP; then since

$$SY : YM :: SA : AX, \quad (\textit{Euclid}, \text{VI. } 2)$$

$$\text{and } SY' : Y'M :: SA' : A'X, \quad (\textit{Euclid}, \text{VI. } 2)$$

$$\therefore\ SY : YM :: SY' : Y'M;$$

or, alternately,

$$SY : SY' :: YM : Y'M,$$

and the angle YPY' in a semicircle is a right angle,

$$\therefore\ PY \text{ bisects the angle } SPM,^*$$

$$\therefore\ SP : PM :: SY : YM,$$

$$:: SA : AX.$$

So we may show that

$$SP' : P'M :: SY : YM,$$

$$:: SA : AX.$$

$\therefore$ P and P' are points on the curve.

---

* For, if not, make the angle YPs equal to YPM; then

$$sY : YM :: sP : PM. \quad (\textit{Euclid}, \text{VI. } 3.)$$

And since, if PY bisect sPM, PY', being at right angles to PY, also bisects the angle sPM',

$$\therefore\ sY' : Y'M :: sP : PM. \quad (\textit{Euclid}, \text{VI. A.})$$

$$\text{Hence } sY : YM :: sY' : Y'M,$$

$$\text{or } sY : sY' :: YM : Y'M,$$

$\therefore$ the points S and s coincide.

In the same way, by taking other points on the directrix, we may obtain as many more points on the curve as we please.

Cor. 1. Since, corresponding to every point P on the curve, there is a point P' situated in precisely the same manner with respect to $A'Y'$ as P is with respect to AY, it is clear that if we make $A'S'$ equal to AS, and $A'X'$ equal to AX, and draw $X'M'$ at right angles to AX', the curve could be equally well described with S' as focus and $M'X'$ as directrix.

The ellipse is therefore symmetrical, not only with respect to the line AA', but also with respect to the line OC drawn through the middle point of YY' at right angles to and bisecting AA'.

Cor. 2. The line OP will bisect the angle SPS'.

Let OP meet SS' in G. Produce MP to meet $X'M'$ in M'. and draw OM' passing through the focus S'; then

$$SP : PM :: S'P : PM',$$

or, alternately, $SP : S'P :: PM : PM'$. (1)

Again, $SG : PM :: S'G : PM'$,

or, alternately, $SG : S'G :: PM : PM'$, (2)

$\therefore$ from (1) and (2)

$$SP : S'P :: SG : S'G,$$

$\therefore$ PG bisects the angle SPS'. (*Euclid*, VI. 3.)

It will be shown hereafter (*Prop.* XI.) that the normal to the ellipse at the point P also bisects the angle SPS'. Hence the ellipse and circle have the same tangent at the point P.

The ellipse will consequently touch all the infinite series of circles which can be described in the same manner as the one in the figure by taking different points on the directrix.

PROP. II.

23. If C be the middle point of AA'. then CA is a mean proportional between CS and CX,

$$\text{or } CS \,.\, CX = CA^2. \quad (\textit{See fig. Prop.}\ \text{III.})$$

$$\text{Since } SA' : A'X :: SA : AX.$$

$$\text{Alternately } SA' : SA :: A'X : AX,$$

$$\therefore\ SA' + SA : SA :: A'X + AX : AX;$$

$$\text{or } AA' : SA :: XX' : AX,$$

$$\therefore\ AA' : XX' :: SA : AX,$$

$$\text{or } CA : CX :: SA : AX. \quad (1)^*$$

$$\text{Again, } SA' : SA :: A'X : AX,$$

$$\therefore\ SA' - SA : SA :: A'X - AX : AX;$$

$$\text{or } SS' : SA :: AA' : AX.$$

$$\text{Alternately } SS' : AA' :: SA : AX;$$

$$\text{or } CS : CA :: SA : AX. \quad (2)$$

Hence from (1) and (2)

$$CA : CX :: CS : CA,$$

$$\therefore\ CA^2 = CX \,.\, CS.$$

or CA is a mean proportional between CS and CX.

COR. Since the three lines CS, CA, CX are proportional, therefore, by the definition of duplicate ratio and *Euclid*, VI. 20 *Cor.*

$$CS : CX : CS^2 : CA^2. \quad (3)$$

PROP. III.

24. If P be any point on the ellipse, then

$$SP + S'P = AA'.$$

---

* N.B. The results (1), (2), (3) should be remembered, as they will frequently be referred to.



$$\begin{aligned}
\text{Since } SP : PM &:: SA : AX,\\
\text{and } SA : AX &:: AA' : XX', \quad (\textit{Prop.}\ \text{II.})\\
\therefore\ SP : PM &:: AA' : XX',\\
\text{So } S'P : PM' &:: AA' : XX',\\
\therefore\ SP + S'P : PM + PM' &:: AA' : XX'.\\
\text{But } PM + PM' &= MM' = XX',\\
\therefore\ SP + S'P &= AA'.
\end{aligned}$$

COR. 1. By means of this property the ellipse may be practically described and the form of the curve determined.

Let a string, equal in length to AA', have its ends fastened to two points S and S'; and let it be kept stretched by means of the point of a pencil at P; then since $SP + S'P$ will be always equal to AA', the point P will trace out the ellipse.

COR. 2. The line AA' is the longest line that can be drawn in the ellipse.

For, if any other line PQ be drawn, then

$$\begin{aligned}
SP + SQ &> PQ,\\
\text{and } S'P + S'Q &> PQ,\\
\therefore\ SP + S'P + SQ + S'Q &> 2PQ,\\
\text{or } AA' &> PQ.
\end{aligned}$$

25. DEF. If BCB' be drawn at right angles to ACA', meeting the ellipse in B and B', it will be seen further on (*Prop.* XIII. *Cor.* 2) that BCB' is the shortest chord that can be drawn through the *centre* of the ellipse. (*See fig. Prop.* IV.)

AA' is called the *Major Axis* and BB' the *Minor Axis* of the ellipse.

In most geometrical treatises the ellipse is defined as the curve traced out by a point which moves in such a manner that the sum of its distances from two fixed points is always the same; but it appears that the properties of the curve are more clearly exhibited by defining it in a manner analogous to the parabola, and deducing *immediately* from that definition the property in question.

Having now shown that one definition necessarily includes the other, we are at liberty in our future investigations to make use of whichever property is most convenient.

Prop. IV.

26. If BC be the semi-minor axis of the ellipse, then

$$BC^2 = CA^2 - CS^2;$$

and if SL be the semi-latus rectum,

$$SL \, . \, AC = BC^2.$$



Join SB, $S'B$; then

$$\text{since } SB + S'B = AA', \quad (\textit{Prop. } \text{III.})$$
$$\text{and that } SB = S'B,$$
$$\therefore \; SB = AC.$$
$$\text{But } BC^2 = SB^2 - CS^2,$$
$$\therefore \; BC^2 = CA^2 - CS^2.$$

Again, $SL : SX :: SA : AX,$

$$:: CS : CA, \quad (\textit{Prop.}\ \text{II.})$$

$$\begin{aligned} \therefore\ SL\,.\,AC &= CS\,.\,SX, \\ &= CS\,.\,CX - CS^2, \quad (\textit{Euclid},\ \text{II. 3,}) \\ &= CA^2 - CS^2, \quad (\textit{Prop.}\ \text{II.}) \\ &= BC^2. \end{aligned}$$

Prop. V.

27. The sum of the distances of *any* point from the foci of an ellipse will be less or greater than AA' according as the point is inside or outside the ellipse.



(1.) Let Q be a point inside the ellipse.

Join SQ, $S'Q$; and produce SQ to meet the ellipse in P; join $S'P$; then

$$\text{since } S'P + QP > S'Q,$$

$$\therefore\ S'P + SP > S'Q + SQ.$$

$$\text{But } S'P + SP = AA', \quad (\textit{Prop.}\ \text{III.})$$

$$\therefore\ SQ + SQ' < AA'.$$

(2.) Let Q be a point outside the ellipse.

Join SQ, $S'Q$, and let SQ meet the ellipse in the point P; join $S'P$; then

$$\text{since } S'Q + QP > S'P,$$

$$\therefore\ S'Q + SQ > SP + S'P,$$

$$\text{But } SP + S'P = AA', \quad (\textit{Prop.}\ \text{III.})$$

$$\therefore\ SQ + S'Q > AA'.$$

Cor. Conversely, a point will be inside or outside the ellipse, according as the sum of its distances from the foci is less or greater than AA'.

28. Def. If a point P' be taken on the ellipse near to P, (*see fig. Prop.* VI.) and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P, is called the *Tangent* to the ellipse at the point P.

Prop. VI.

If the tangent to the ellipse at any point P intersect the directrix in the point Z, and if S be the focus corresponding to the directrix on which Z is situated, then SZ will be at right angles to SP.



Let P' be a point on the ellipse near to P.

Draw the chord PP', and produce it to meet the directrix in Z; join SZ.

Draw PM, $P'M'$ at right angles to the directrix, and join SP, SP'.

Produce PS to meet the ellipse in the point Q; then since the triangles ZMP, $ZM'P'$ are similar,

$$\therefore ZP : ZP' :: MP : M'P',$$
$$:: SP : SP',$$

$\therefore$ SZ bisects the angle $P'SQ$. (*Euclid*, VI. *Prop.* A.)

Now when P' is indefinitely near to P, and PP' becomes the tangent at the point P, the angle PSP' becomes indefinitely small, while the angle QSP' approaches two right angles; and therefore the angle ZSP' being half of the angle $P'SQ$, becomes ultimately a right angle.

Hence when PZ becomes the tangent at the point P,

the angle ZSP is a right angle,

or SZ is perpendicular to SP.

Cor. 1. Conversely, if SZ be drawn at right angles to SP, meeting the directrix in Z, and PZ be joined, PZ will be the tangent at P.

Cor. 2. If ZP be produced to meet the other directrix on the point Z' and $S'Z'$ be joined, then

$S'Z'$ is at right angles to $S'P$.

Cor. 3. The tangents at the extremities of the latus rectum or double ordinate through the focus meet the axis produced in the point X.

Prop. VII.

The tangent to the ellipse at any point P makes equal angles with the focal distances SP and $S'P$.

Let the tangent at P meet the directrices in Z and Z'.

Draw MPM' at right angles to the directrices, meeting them in M and M' respectively; join SZ, $S'Z'$; then

$$SP : PM :: S'P : PM';$$

and since the triangles MPZ, $M'PZ'$, are similar,

$$PM : PZ :: PM' : PZ',$$
$$\therefore SP : PZ :: S'P : PZ'. \quad (\textit{Ex æquali.})$$



Now in the triangles SPZ, $S'PZ'$, because the sides about the angles SPZ, $S'PZ'$ are proportional, and the angles PSZ, $PS'Z'$ are equal, being right angles, and the angles SZP, $S'Z'P$ are each less than a right angle,

$\therefore$ the triangles SPZ and $S'PZ'$ are similar, (*Euclid*, VI. 7)

$$\therefore \text{ the angle } SPZ = \text{the angle } S'PZ'.$$

COR. If $S'P$ be produced to W; then

$$\text{the angle } SPZ = \text{the angle } WPZ.$$

PROP. VIII.

The tangents at the extremities of a focal chord intersect in the directrix.

Let PSQ be a focal chord, and let the tangent at P meet the directrix in Z.

Join SZ; then

the angle ZSP is a right angle, (*Prop.* VI.)
and $\therefore$ also the angle ZSQ is a right angle,
$\therefore$ ZQ is the tangent at Q; (*Prop.* VI. *Cor.* 1)
or the tangents at the extremities of a focal chord intersect in the directrix.

PROP. IX.

29. If the tangent at P meet the axis major produced in T, and PN be the ordinate of the point P, then

$$CT \,.\, CN = CA^2.$$



Draw MPM' parallel to the axis major meeting the directrices in M and M'; and produce $S'P$ to W; then, since PT bisects the angle SPW, (*Prop.* VII. *Cor.*)

$$\begin{aligned} \therefore\ S'T : ST &:: S'P : SP, \quad (\textit{Euclid}, \text{VI. A.}) \\ &:: PM' : PM, \\ &:: X'N : XN, \end{aligned}$$

$$\therefore\ S'T + ST : S'T - ST :: X'N + XN : X'N - XN;$$

$$\text{or } 2CT : 2CS :: 2CX : 2CN,$$

$$\text{or } CT : CS :: CX : CN,$$

$$\begin{aligned} \therefore\ CT \,.\, CN &= CS \,.\, CX, \\ &= CA^2. \quad (\textit{Prop}\ \text{II.}) \end{aligned}$$

PROP. X.

If on the major axis of an ellipse as diameter a circle be described and a common ordinate NPQ be drawn meeting the ellipse in P and the circle in Q, then the tangents to the ellipse and circle respectively at the points P and Q will meet the major axis produced in the same point.



Let the tangent to the ellipse at P meet the major axis produced in T; join CQ, QT; then, by the last Proposition,

$$CT \,.\, CN = CA^2 = CQ^2,$$

$\therefore$ the angle CQT is a right angle.

And therefore QT is the tangent to the circle at Q; or the tangents at P and Q meet the major axis produced in the same point T.

The circle described on AA' as diameter is called the *Auxiliary Circle* on account of the important aid that it affords in investigating the properties of the ellipse.

30. DEF. The line PG, drawn at right angles to the tangent PT, is called the *Normal* to the ellipse at the point P.

PROP. XI.

If the normal at P meet the major axis in the point G; then

$$SG : SP :: CS : CA,$$



Since PG is at right angles to TPt,
$\therefore$ the angle $GPT =$ the angle GPt.
But the angle $SPT =$ the angle $S'Pt$, (*Prop.* VII.)
$\therefore$ the angle $SPG =$ the angle $S'PG$,
or PG bisects the angle SPS',

$$\therefore SG : S'G :: SP : S'P, \quad (\textit{Euclid}, \text{VI. } 3.)$$

$$\therefore SG : SG + S'G :: SP : SP + S'P;$$

$$\text{or } SG : SS' :: SP : AA';$$

$$\text{or } SG : SP :: SS' : AA';$$

$$\text{or } SG : SP :: CS : CA.$$

Cor. Hence also,

$$S'G : S'P :: CS : CA.$$

Prop. XII.

31. If the normal at P meet the major axis in G, and PN be the ordinate at the point P, then (*see fig. Prop.* XI.)

$$NG : NC :: BC^2 : AC^2.$$

Draw MPM' parallel to the axis meeting the directrices in M and M'; join SP, $S'P$; then, since PG bisects the angle SPS', (*Prop.* XI.)

$$\begin{aligned} \therefore S'G : SG &:: S'P : SP, \\ &:: PM' : PM, \\ &:: X'N : XN, \end{aligned}$$

$$\therefore S'G - SG : S'G + SG :: X'N - XN : X'N + XN;$$

$$\text{or } 2\,CG : SS' :: 2\,CN : XX'.$$

$$\text{Alternately, } 2CG : 2CN :: SS' : XX';$$

$$\text{or } CG : CN :: CS : CX,$$

$$:: CS^2 : CA^2, \text{ (Prop. II. Cor.)}$$

$$\therefore\ CN - CG : CN :: CA^2 - CS^2 : CA^2;$$

$$\text{or } NG : CN :: BC^2 : AC^2.$$

PROP. XIII.

32. If PN be the ordinate of any point P on the ellipse then

$$PN^2 : AN.A'N :: BC^2 : AC^2.$$



Produce NP to meet the auxiliary circle in the point Q, and draw the tangents PT, QT meeting the major axis produced in the point T. (*Prop.* X.)

Join CQ, and let the normal at P meet the ellipse in G; then, by the last Proposition,

$$NG : CN :: BC^2 : AC^2.$$

And rectangles of the same altitude are to another as their bases,

$$\therefore\ TN . NG : TN . CN :: BC^2 : AC^2;$$

$$\text{or } PN^2 : QN^2 :: BC^2 : AC^2. \text{ (Euclid, VI. 8, Cor.)}$$

$$\text{But } QN^2 = AN . A'N,$$

since the angle AQA' in a semicircle is a right angle,

$$\therefore\ PN^2 : AN . A'N :: BC^2 : AC^2.$$

COR. 1. Also $PN : QN :: BC : AC$.

This result is the basis of many of the future Propositions of the ellipse.

Cor. 2. Since $PN^2 : QN^2 :: BC^2 : AC^2$,

$$\therefore PN^2 : AC^2 - CN^2 :: BC^2 : AC^2,$$

$$\therefore PN^2 : AC^2 - CN^2 - PN^2 :: BC^2 : AC^2 - BC^2,$$

$$\text{or } PN^2 : AC^2 - CP^2 :: BC^2 : AC^2 - BC^2.$$

Now PN^2 is always less than BC^2,

$\therefore$ CP^2 is greater than BC^2,

$\therefore$ BC is the shortest line that can be drawn to the ellipse from the centre.

Prop. XIV.

If the tangent at any point P of an ellipse meet the minor axis CB produced in t, and Pn be drawn at right angles to CB; then

$$Ct \quad Cn = BC^2$$



Draw the common ordinate NPQ to the ellipse and the auxiliary circle; and let the tangents at P and Q to the ellipse and circle respectively meet the major axis produced in T (*Prop.* X.) and the minor axis produced in t and U.

Join CQ meeting Pn in R; then since PR is parallel to CN,

$$CR : CQ :: PN : QN,$$

$$:: BC : AC. \quad (\textit{Prop.}\ \text{XIII. } \textit{Cor.}\ 1.)$$

$$\text{But } CQ = AC,$$
$$\therefore\ CR = BC. \quad (1)$$

Again, joining Rt,

$$Ct : CU :: PN : QN,$$
$$:: CR : CQ,$$
$$\therefore\ Rt \text{ is parallel to } QU,$$
$$\therefore\ CRt \text{ is a right angle,}$$
$$\therefore\ Ct \quad Cn = CR^2 \ (\textit{Euclid}, \text{ VI. 8, } \textit{Cor.})$$
$$\text{But } CR = BC, \quad (1)$$
$$\therefore\ Ct \,.\, Cn = BC^2.$$

This proposition also admits of a demonstration similar to that given for the corresponding property of the hyperbola.

Prop. XV.

33. If from the foci S and S', SY, and $S'Y'$ are drawn at right angles to the tangent at P, then Y and Y' are on the circumference of the auxiliary circle, and

$$SY \,.\, S'Y' = BC^2.$$



Join SP, $S'P$, and produce SY and $S'P$ to meet in W; and join CY; then

since the angle SPY = the angle WPY, (*Prop.* VII. *Cor.*)
and the angle SYP = the angle WYP,
and the side PY is common to the triangles SPY, WPY,
$\therefore$ the triangle SPY = the triangle WPY in all respects,

$$\therefore SP = PW,$$
$$\therefore SP + S'P = S'W.$$
$$\text{But } SP + S'P = AA', \quad (\textit{Prop.}\ \text{III.})$$
$$\therefore S'W = AA'.$$

Again $\because$ $SC = S'C$ and $SY = YW$,

$$\therefore SC : S'C :: SY : YW,$$
$$\therefore CY \text{ is parallel to } S'W,$$
$$\therefore CY : S'W :: CS : SS',$$
$$\therefore CY = \tfrac{1}{2}S'W = CA.$$
$$\text{So } CY' = CA,$$

$\therefore$ Y and Y' are points on the auxiliary circle.

Next let YS be produced to meet the auxilliary circle in Z, and join ZY'; then

since the angle ZYY' is a right angle,
$\therefore$ ZY' passes through the centre C,
$\therefore$ the angle SCZ = the angle $S'CY'$.

$$\therefore SZ = S'Y',$$
$$\begin{aligned} \therefore SY \,.\, S'Y' &= SY \,.\, SZ, \\ &= AS \,.\, A'S, \quad (\textit{Euclid}, \text{III. } 35) \\ &= CA^2 - CS^2, \quad (\textit{Euclid}, \text{II. } 5) \\ &= BC^2. \quad (\textit{Prop.}\ \text{IV.}) \end{aligned}$$

Cor. If CD be drawn parallel to the tangent at P, meeting $S'P$ in E; then since the figure $CYPE$ is a parallelogram,

$$\therefore\ PE = CY = AC.$$

Prop. XVI.

34. To draw a pair of tangents to an ellipse from an external point O.



With centre S' and radius equal to AA' describe a circle.

Join OS, OS'; and let SO or $S'O$ produced meet the circle in the point I.

Now, if O be a point outside the circle MIM', it is evident that OS is greater than OI; and if O be inside the circle,

$$\text{since } OS + OS' > AA' \text{ or } S'I, \quad (\textit{Prop.}\ \text{V.})$$
$$\therefore\ OS > OI.$$

With centre O and radius OS describe another circle cutting the former in the points M and M', which it will always do since OS is greater than OI.

Join $S'M$, $S'M'$, meeting the ellipse in the points P and P'.

Join OP, OP'; these will be the tangents required.

Join SP, SP'; then, since

$$SP + S'P = AA' = S'M,$$

$$\therefore\ SP = PM.$$

And $\because$ SP, $PO = MP$, PO each to each,

and $OS = OM$,

$\therefore$ the angle OPS = the angle OPM,

$\therefore$ OP is the tangent at P. (*Prop.* VII. *Cor.*)

So OP' is the tangent at P'.

PROP. XVII.

If from a point O a pair of tangents OP, OP' be drawn to an ellipse, then OP and OP' will subtend equal angles at either focus.

Join SP, $S'P$; SP', $S'P'$; and produce $S'P$, $S'P'$ to M and M', making PM equal to SP, and $P'M'$ equal to SP.

Join OM, OM'; OS, OS'.

Then since OP, $PS = OP$, PM, each to each,

and the angle OPS = the angle OPM, (*Prop.* VII. *Cor.*)

$$\therefore\ OS = OM,$$

and the angle OSP = the angle OMP.

So $OS = OM'$,

and the angle OSP' = the angle $OM'P'$,

$$\therefore\ OM = OM'.$$

Again, $\because$ $S'M = S'P + SP = AA'$,

and $S'M' = S'P' + SP' = AA'$,

$$\therefore\ S'M = S'M',$$

And $\because$ OS', $S'M = OS'$, $S'M'$, each to each,

and $OM = OM'$,

$\therefore$ the angle $OS'M$ = the angle $OS'M'$,

and the angle OMS' = the angle $OM'S'$.

But the angle OMS' = the angle OSP,
and the angle $OM'S'$ = the angle OSP',
∴ the angle OSP = the angle OSP',
∴ OP and OP' subtend equal angles at either focus.

Prop. XVIII.

35. If from an external point O a pair of tangents OQ, OQ' be drawn to an ellipse, and CO be joined meeting the chord QQ' in V, and the ellipse in P; then

(1.) QQ' will be bisected in V.
(2.) The tangent at P will be parallel to QQ'.
(3.) CP will be a mean proportional between CV and CO.



Produce OQ, OQ' to meet the major axis produced in T and T'.

Draw the ordinates NQ, $N'Q'$, and produce them to meet the circle in q and q'.

Then Tq and $T'q'$ will be tangents to the auxiliary circle. (*Prop.* X.)

Let Tq and $T'q'$ be produced to meet in o; join Co meeting the chord qq' in v, and the circle in p.

Now, since the corresponding ordinates of the ellipse and auxiliary circle are in the constant ratio of BC to AC, the three lines ol, pm, vn drawn at right angles to AA' will pass through the points O, P, V respectively.

For, according as O is the point where ol meets TQ or $T'Q'$ we shall have

$$\begin{aligned} lO : lo &:: NQ : Nq, \\ &:: BC : AC; \\ \text{or } lO : lo &:: N'Q' : N'q', \\ &:: BC : AC, \end{aligned}$$

$\therefore Oo$ is perpendicular to AA'.

So Pp and Vv are perpendicular to AA',

$\therefore Oo, Pp, Vv$ are parallel.

Hence (1.) $QV : VQ' :: qv : vq'$.

But $qv = vq'$ from the circle,

$\therefore QV = VQ'$;

or QQ' is bisected in V.

(2.) Since $NQ : Nq :: N'Q' : N'q'$

it is evident that QQ' and qq' will meet the axis produced in the same point.

Also the tangents to the ellipse and circle at P and p respectively will meet the axis in the same point.

Now in the circle the tangent at p is manifestly parallel to qq',

and $NQ : Nq :: mP : mp$,

$\therefore$ the tangent at P is parallel to QQ'.

(3.) If Cq be joined, since the angle Cqo is a right angle and Co is perpendicular to qq',

$\therefore Cv : Cq :: Cq : Co$, (*Euclid*, VI. 8 *Cor.*)

or, since $Cq = Cp$,

$Cv : Cp :: Cp : Co$.

$$\begin{aligned} \text{But } Cv : Cp &:: CV : CP, \\ \text{and } Cp : Co &:: CP : CO, \\ \therefore CV : CP &:: CP : CO, \\ \therefore CO \, . \, CV &= CP^2. \end{aligned}$$

Cor. From this it is manifest that if any number of chords be drawn parallel to each other in an ellipse, their middle points will all lie on the line drawn from the centre to the point where the tangent parallel to the chord meets the ellipse.

Def. The line PCP' drawn through the centre of an ellipse and meeting the curve in P and P', is called a *Diameter.*

The diameter consequently bisects all chords parallel to the tangents at its extremities; and the tangents at the extremities of any chord will intersect the diameter corresponding to that chord in the same point.

36. Def. If CD be drawn parallel to the tangent at P, then CD is said to be *conjugate* to CP.

Prop. XIX.

In the ellipse if CD be conjugate to CP, then will CP be conjugate to CD.



Draw the ordinates PN, DR, and produce them to meet the auxiliary circle in the points p, d.

Join CP, Cp; CD, Cd; and draw the tangents TP, Tp; $T'D$, $T'd$.

Now, since CD is parallel to PT,

$\therefore$ the triangle PNT is similar to the triangle DRC.

$\therefore TN : CR :: PN : DR,$

$:: Np : Rd,$ (*Prop.* XIII. *Cor.*)

$\therefore Tp$ is parallel to Cd,

$\therefore$ the angle pCd is a right angle,

$\therefore Cp$ is parallel to $T'd$,

$\therefore$ the triangle pCN is similar to the triangle $dT'R$,

$\therefore NC : RT' :: Np : Rd,$

$:: NP : RD,$

$\therefore CP$ is parallel to DT',

$\therefore CP$ is conjugate to CD.

Cor. Since CRd and CNp are each similar to dRT', *Euclid*, VI. 8)

$\therefore$ the triangle CRd is similar to the triangle CNp,

and the side $Cd =$ the side Cp,

$\therefore$ the triangle $CRd =$ the triangle CNp in all respects,

$\therefore CN = Rd$, and $CR = Np$.

Hence $DR : CN :: DR : Rd,$

$:: BC : AC;$

also $PN : CR :: PN : Np,$

$:: BC : AC.$

Prop. XX.

37. If CP and CD be conjugate semi-diameters, and PN, DR be the ordinates of the points P and D; then

(1.) $CN^2 + CR^2 = AC^2$.

(2.) $PN^2 + DR^2 = BC^2$.

(3.) $CP^2 + CD^2 = AC^2 + BC^2$.



Produce NP, RD to meet the auxiliary circle in the points p, d; then

$$CN = Rd, \quad (\textit{Prop.}\ \text{XIX.}\ \textit{Cor.})$$

$$\therefore CN^2 + CR^2 = Rd^2 + CR^2,$$

$$= Cd^2,$$

$$= CA^2.$$

$$\text{Again, } PN : Np :: BC : AC,$$

$$\therefore PN^2 : Np^2 :: BC^2 : AC^2.$$

$$\text{So } DR^2 : Rd^2 :: BC^2 : AC^2,$$

$$\therefore PN^2 + DR^2 : Np^2 + Rd^2 :: BC^2 : AC^2;$$

$$\text{but } NP^2 + Rd^2 = CR^2 + CN^2,$$

$$= AC^2,$$

$$\therefore PN^2 + DR^2 = BC^2,$$

$$\text{and } CN^2 + CR^2 = AC^2,$$

$$\therefore CP^2 + CD^2 = AC^2 + BC^2.$$

38. Def. A line QV drawn parallel to the tangent at P, and meeting CP in V, is called an *Ordinate* to the diameter CP.

Prop. XXI.

If QV be any ordinate to the diameter PCP', and CD be conjugate to CP; then

$$QV^2 : PV \,.\, P'V :: CD^2 : CP^2.$$

Draw the tangent UQW meeting CP and CD produced in U and W; and draw QR parallel to CP, meeting CD in R.

Now, since $CR : CD :: CD : CW$, (*Prop.* XVIII.)

$\therefore CR^2 : CD^2 :: CR : CW$, (*Euclid*, VI. 20 *Cor.*)

or $QV^2 : CD^2 :: UV : CU$.



Again,

since $CU : CP :: CP : CV$, (*Prop.* XVIII.)

$\therefore CU : CV :: CP^2 : CV^2$, (*Euclid*, VI. 20 *Cor.*)

$\therefore CU - CV : CU :: CP^2 - CV^2 : CP^2$,

or $UV : CU :: PV \,.\, P'V : CP^2$.

Hence $QV^2 : CD^2 :: PV \,.\, P'V : CP^2$.

or $QV^2 : PV \,.\, P'V :: CD^2 : CP^2$.

Prop. XXII.

39. The area of any parallelogram formed by drawing tangents to an ellipse at the extremities of a pair of conjugate

diameters is equal to the rectangle contained by the axes of the ellipse.

Let PCP', DCD' be a pair of conjugate diameters, and let a parallelogram be formed by drawing tangents at the points P, P', D, D'.



Let the tangent at P meet CA produced in T; join $D'T$.

Draw the ordinates PN, DR, $D'R'$; then since PT is parallel to CD', the parallelogram PD' is double the triangle CTD', and therefore equal to the rectangle contained by CT and $D'R'$.

Now $D'R' : CN :: BC : AC$, (*Prop.* XIX. *Cor.*)

$\therefore CT \cdot D'R' : CT \cdot CN :: BC : AC$, (*Euclid*, VI. 1.)

$:: BC \cdot AC : AC^2$. (*Euclid*, VI. 1.)

But $CT \cdot CN = AC^2$,

$\therefore CT \cdot D'R' = AC \cdot BC$,

$\therefore$ the parallelogram $LL' = 4$ the parallelogram PD',

$= 4\, AC \cdot BC$,

$= AA' \cdot BB$.

Cor. If PF be drawn at right angles to DCD' meeting CD' in F; then

$$\begin{aligned} PF \,.\, CD' &= \text{area of parallelogram } PD', \\ &= AC \,.\, BC. \end{aligned}$$

Prop. XXIII.

40. If CP and CD be conjugate diameters, and PF be drawn at right angles to CD meeting CA in G, then

$$PF \,.\, PG = BC^2.$$



Draw the ordinate PN, and produce it to meet CD' in K.

Also draw Pn at right angles to CB, and let the tangent at P meet CB produced in t.

Now since the angles at N and F are right angles, it is evident that a circle may be described about the quadrilateral figure $NKFG$;

$$\begin{aligned} \therefore PG \,.\, PF &= PN \,.\, PK, \quad (\textit{Euclid}, \text{ III. 36 } \textit{Cor.}) \\ &= Ct \,.\, Cn, \\ &= BC^2. \quad (\textit{Prop.} \text{ XIV.}) \end{aligned}$$

PROP. XXIV.

41. If P be any point on the ellipse, and CD be conjugate to CP, then

$$SP \,.\, S'P = CD^2.$$



Draw the normal PG and produce it to meet CD' in F; then since CD' is parallel to the tangent at P,

$$\therefore\ PF \text{ is at right angles to } CD',$$

$$\therefore\ PF \,.\, CD = AC \,.\, BC, \quad (\textit{Prop.}\ \text{XXII. } \textit{Cor.})$$

$$\text{and } PF \,.\, PG = BC^2 = BC \,.\, BC, \quad (\textit{Prop.}\ \text{XXIII.})$$

$$\therefore\ CD : PG :: AC : BC. \quad (1)$$

$$\text{Again, } SP : SG :: CA : CS, \quad (\textit{Prop.}\ \text{XI.})$$

$$S'P : S'G :: CA : CS. \quad (\textit{Prop.}\ \text{XI.})$$

$$\text{Compounding } SP \,.\, S'P : SG \,.\, S'G :: CA^2 : CS^2,$$

$$\therefore\ SP \,.\, S'P : SP \,.\, S'P - SG \,.\, S'G :: CA^2 : CA^2 - CS^2.$$

$$\text{But } SP \,.\, S'P - SG \,.\, S'G = PG^2, \quad (\textit{Euclid}, \text{ VI. } \textit{Prop.}\ \text{B})$$

$$\therefore\ SP \,.\, S'P : PG^2 :: CA^2 : BC^2.$$

$$\text{But from (1) } CD^2 : PG^2 :: CA^2 : BC^2,$$

$$\therefore\ SP \,.\, S'P = CD^2.$$

This proposition may also be very easily deduced from *Prop.* XV.

PROP. XXV.

42. The area of the ellipse is to the area of the auxiliary circle as BC to AC.



Let PN and $P'N'$ be two ordinates of the ellipse near together.

Produce NP, $N'P'$, to meet the auxiliary circle in Q and Q'.

Draw Pm, Qn, perpendicular to $Q'N'$.

Then

the parallelogram PN' : the parallelogram $QN' :: PN : QN$,
$:: BC : AC$.

And the same will be true for all the parallelograms that can be similarly described in the ellipse and auxiliary circle.

Hence the sum of all the parallelograms inscribed in the ellipse is to the sum of all the parallelograms inscribed in the circle as BC to AC.

And this holds however the number of parallelograms be increased.

But when the number of parallelograms is increased, and the breadth of each diminished indefinitely, the sum of the parallelograms inscribed in the ellipse will be equal to the area of the ellipse, and the sum of those inscribed in the circle to the area of the circle. Hence

the area of the ellipse : the area of the circle $:: BC : AC$.

43. Def. If with a point O on the normal at P as centre, and OP as radius, a circle be described touching the ellipse at P, and cutting it in Q; then, when the point Q is made to approach indefinitely near to P, the circle is called the *Circle of Curvature* at the point P.

Prop. XXVI.

If PH be the chord of the circle of curvature at the point P of an ellipse, which passes through the centre; then

$$PH \, . \, CP = 2CD^2.$$

Let PT be the tangent, and PG the normal at the point P.



With centre O, and radius OP, describe a circle cutting the ellipse in the point Q.

Draw RQW parallel to CP, meeting the circle in W, and TP produced in R.

Also draw QV parallel to PR, meeting the diameter PP' in V; then since RP touches the circle at P,

$$\therefore RQ \,.\, RW = PR^2, \text{ (\textit{Euclid}, III. 36)}$$

$$\text{or } PV \,.\, RW = QV^2.$$

$$\text{But } QV^2 : PV \,.\, P'V :: CD^2 : CP^2, \text{ (\textit{Prop.} XXI.)}$$

$$\therefore PV \,.\, RW : PV \,.\, P'V :: CD^2 : CP^2,$$

$$\text{or } RW : P'V :: CD^2 : CP^2.$$

Now, when the circle becomes the circle of curvature at P, the points R and Q move up to, and coincide with P, and the lines RW and PH become equal, while

$P'V$ becomes equal to PP', or $2CP$.

Hence, $PH : 2CP :: CD^2 : CP^2$,

$$\therefore PH \,.\, CP : 2CP^2 :: 2CD^2 : 2CP^2,$$

$$\therefore PH \,.\, CP = 2CD^2.$$

Prop. XXVII.

If PU be the diameter of the circle of curvature at the point P of the ellipse, and PF be drawn at right angles to CD; then

$$PU \,.\, PF = 2CD^2.$$

Since the triangle PHU is similar to the triangle PFC,

$$\therefore PU : PH :: CP : PF,$$

$$\therefore PU \,.\, PF = PH \,.\, CP,$$

$$= 2CD^2. \quad \text{(\textit{Prop.} XXVI.)}$$

Prop. XXVIII.

If PI be the chord of the circle of curvature through the focus of the ellipse; then

$$PI \,.\, AC = 2CD^2.$$

Let PI meet CD in E; then, since the triangles PIU and PEF are similar,

$$\therefore\ PI : PU :: PF : PE.$$

But $PE = AC$, (*Prop.* XV. *Cor.*)

$$\therefore\ PI : PU :: PF : AC,$$

$$\therefore\ PI \,.\, AC = PU \,.\, PF,$$

$$= 2CD^2. \quad (\textit{Prop.}\ \text{XXVII.})$$

Prop. XXIX.

44. If two chords of an ellipse intersect one another, the rectangles contained by their segments are proportional to the squares of the diameters parallel to them.

Let POP' be any chord drawn through the point O, and let CD be the semi-diameter parallel to it.

Draw the ordinates NP, $N'P'$, MD, and produce them to meet the auxiliary circle in Q, Q', D'; then

since $NP : NQ :: N'P' : N'Q'$, (*Prop.* XIII. *Cor.*)

it is evident that PP' and QQ' will meet the axis produced in the same point T.



Also since $NP : NQ :: MD : MD'$, (*Prop.* XIII. *Cor.*)

and TPP' is parallel to CD,

$\therefore$ TQQ' is parallel to CD'.

Draw EO parallel to NQ or $N'Q'$, and produce it to meet QQ' in O'; then

$$PO : QO' :: TO : TO',$$
$$\text{and } P'O : Q'O' :: TO : TO',$$
$$\therefore PO \,.\, P'O : QO' \,.\, Q'O' :: TO^2 : TO'^2,$$
$$:: CD^2 : CD'^2,$$
$$:: CD^2 : AC^2.$$

Alternately, $PO \,.\, P'O : CD^2 :: QO' \,.\, Q'O' : AC^2$.

Again, if through the point O any other chord pOp' be drawn,

$$\text{since } EO : EO' :: BC : AC,$$

it is manifest that the corresponding chord qq' in the auxiliary circle will pass through the point O'; and if Cd be the semi-diameter parallel to pp' we shall have as before,

$$pO \,.\, p'O : Cd^2 :: qO' \,.\, q'O' : AC^2.$$

But $QO' \,.\, Q'O' = qO' \,.\, q'O'$, (*Euclid*, III. 35)

$$\therefore PO \,.\, P'O : CD^2 :: pO \,.\, p'O : Cd^2.$$
$$\text{or } PO \,.\, P'O : pO \,.\, p'O :: CD^2 : Cd^2.$$

The same result may be shown to be true when the point O is without the ellipse.

Prop. XXX.

If QVQ' be any ordinate to the diameter CP, the circle described through the three points P, Q, Q' will intersect the ellipse in a fourth point, which depends only upon the position of P.

Draw the ordinate PN, and produce it to meet the ellipse in P'; then since, if NT be the subtangent of either P or P',

$$CT \,.\, CN = AC^2, \text{ (*Prop.* IX.)}$$

therefore the tangents at P and P' will meet the major axis produced in the same point T.



Draw PR parallel to TP', meeting the ellipse in R, and QQ' in O; then if CD and CD' be drawn parallel respectively to TP and TP', meeting the ellipse in D and D',

$$PO \,.\, OR : QO \,.\, OQ' :: CD'^2 : CD^2. \quad (\textit{Prop.}\ \text{XXIX.})$$

$$\text{But } CD' = CD, \text{ since } CP' = CP,$$

$$\therefore PO \,.\, OR = QO \,.\, OQ'.$$

Hence, by the converse of *Euclid* III. *Prop.* 22, the point R is on the circle which passes through P, Q, Q'.

Cor. When the point V is brought indefinitely near to P, QQ' coincides with the tangent to the ellipse at P, and becomes also a tangent to the circle at P since Q and Q' are indefinitely near to each other. The circle therefore becomes the circle of curvature at the point P.

Hence, if PR be drawn parallel to the tangent at P', or be equally inclined to the axis with PT, it will meet the ellipse in the point where the circle of curvature at P intersects the ellipse.

PROBLEMS ON THE ELLIPSE.

1. In what position of P is the angle SPS' greatest?

2. The latus rectum is a third proportional to the axis major and axis minor.

3. Construct on the axis minor as base, a rectangle which shall be to the triangle SLS' in the duplicate ratio of the major axis to the minor axis, L being the extremity of the latus rectum.

4. If a series of ellipses be described having the same major axis; the tangents at the extremities of their latera recta will all meet the minor axis in the same point.

5. Find the locus of the centres of all the ellipses having the same focus, and their major axes of the same length, and touching a given straight line.

6. Given the foci, it is required to describe an ellipse touching a given straight line.

7. If PT be a tangent to an ellipse, meeting the axis in T, and AP, $A'P$, be produced to meet the perpendicular to the major axis through T in Q and Q', then $QT = Q'T$.

8. If the angle SBS' be a right angle, prove that $CA^2 = 2CB^2$.

9. If CP be a semi-diameter, and AQO be drawn parallel to CP meeting the curve in Q, and CB produced in O, then $CP^2 = AO \cdot AQ$.

10. If AB, CD, which are not parallel, make equal angles with either axis, the lines AC, BD, as also AD, BC, will make equal angles with either axis.

11. PSp is any focal chord. PA and pA are produced to meet the directrix in Q and q. Prove that the angle QSq is a right angle.

12. If a circle be described touching the axis major in one focus, and passing through one extremity of the axis minor; AC will be a mean proportional between the diameter of this circle and BC.

13. If $PQQ'P'$ be a chord of the auxiliary circle, and a circle be described on the minor axis as diameter, cutting the chord in Q and Q', then $PQ \,.\, P'Q = CS^2$.

14. If PG be the normal at P, and GL be drawn at right angles to SP, then $PL = \frac{1}{2}$ latus rectum.

15. The sum of the squares of the normals at the extremities of conjugate diameters is constant.

16. If on the normal at P, PQ be taken equal to the semi-conjugate diameter CD, the locus of Q is a circle whose radius is $AC - BC$.

17. Find the locus of the intersection of a pair of tangents at right angles to each other.

18. P is any point on an ellipse. To any point Q on the curve draw AQ, $A'Q$, meeting NP in R and S, and prove that $NR \,.\, NS = NP^2$.

19. If PG be a normal, and GL perpendicular to SP, the ratio of GL to PN is constant.

20. If NP produced meet the tangent at the extremity of the latus rectum in Q, then $QN = PS$.

21. In an ellipse the tangent at any point makes a greater angle with the focal distance than with the perpendicular on the directrix.

22. A diameter of an ellipse, parallel to the tangent at any point, meets the focal distances of the point, and from the points of intersection lines are drawn perpendicular to the focal distances. Prove that these lines intersect in the axis minor.

23. The subnormal is a third proportional to CT and BC.

24. If PN be the ordinate of P, prove that $NY : NY' :: PY : PY'$. (*See fig. Prop.* XV.)

25. If from C lines be drawn parallel and perpendicular to the tangent at P, they inclose a part of one of the focal distances of that point equal to the other.

26. If P be a fixed point on an ellipse, and QQ' an ordinate to CP, the circle QPQ' will meet the ellipse in a fixed point.

27. P is any point on an ellipse. Draw PP' parallel to the axis major, and through P' draw $P'Q$, $P'Q'$, making equal angles with the major axis. Join QQ'; then QQ' is parallel to the tangent at P.

28. What parallelogram circumscribing an ellipse has the least area?

29. When is the square of the sum of conjugate diameters least?

30. Given the axes of an ellipse, and the position of one focus, and of one point in the curve, give a geometrical construction for finding the centre.

31. If lines drawn through any point of an ellipse to the extremities of any diameter meet the conjugate CD in M and N, then $CM \cdot CN = CD^2$.

32. If CP and CD be conjugate, prove that

$$(SP - AC)^2 + (SD - AC)^2 = SC^2.$$

33. If CP and CD be conjugate, and BP, BD be joined, as also AD, $A'P$, these latter meeting in O, then $BDOP$ is a parallelogram. When is the area greatest?

34. If PSp, QCq be two parallel chords through the focus and centre of an ellipse, prove that

$$SP \cdot Sp : CQ \cdot Cq :: BC^2 : AC^2.$$

35. If the tangent at the vertex A cut any two conjugate diameters in T and t, then $AT \cdot At = BC^2$.

36. If the tangents at three points P, Q, R, intersect in $R_{\prime}$, $Q_{\prime}$, $P_{\prime}$, prove that

$$PR_{\prime} \cdot P_{\prime}Q \cdot Q_{\prime}R = PQ_{\prime} \cdot R_{\prime}Q \cdot P_{\prime}R.$$

37. If a circle be described touching SP, $S'P$ produced, and the major axis of the ellipse, find the locus of the centre.

38. If from the extremities of the axes of an ellipse any four parallel lines be drawn, the points in which they cut the curve are the extremities of conjugate diameters.

39. If two equal and similar ellipses have a common centre, the points of intersection are at the extremities of diameters at right angles to one another.

40. If PSQ be a focal chord, and X the foot of the directrix, XP and XQ are equally inclined to the axis.

41. OP, OQ are tangents to an ellipse, and PQ is produced to meet the directrices in R, R', prove that

$$RP \,.\, R'P : RQ : R'Q :: OP^2 : OQ^2.$$

42. NPQ is a common ordinate to the ellipse and auxiliary circle. PR, QR are normals at P and Q intersecting in R. The locus of R is a circle whose radius is $AC + BC$.

43. If the conjugate to CP meet SP, $S'P$, or these produced in E, E'; then $SE = S'E'$, and the circles circumscribing SCE, $S'CE'$ are equal.

44. The locus of the middle points of all focal chords in an ellipse is a similar ellipse.

45. The circle described about the triangle SBS' will cut the minor axis in the centre of the circle of curvature at B.

46. The locus of the centre of the circle inscribed in the triangle SPS' is an ellipse.

47. If a circle be described intersecting an ellipse in four points, and chords be drawn through the points of intersection, diameters parallel to the chords will be equal.

48. An ellipse slides between two lines at right angles to each other, find the locus of its centre.

49. If from the focus S perpendiculars be drawn upon the conjugate diameters CP, CD, these perpendiculars produced backward will intersect CD and CP in the directrix.

50. Find the point at which the diameter of curvature is a mean proportional between the major and minor axes.

51. The circle of curvature at a point, where the conjugate diameters are equal, meets the ellipse again at the extremity of the diameter.

52. The locus of the intersection of lines drawn from A, A' at right angles to AP, $A'P$ is an ellipse.

53. If a quadrilateral figure be inscribable in two ellipses whose major axes are parallel or perpendicular, any two of its opposite angles will be equal to two right angles.

54. If CN, NP are the abscissa and ordinate of a point P on a circle whose centre is C, and NQ be taken equal to NP, and be inclined to it at a constant angle, the locus of Q is an ellipse.

55. If two ellipses having the same major axes can be inscribed in a parallelogram, the foci will be on the corners of an equiangular parallelogram.

56. Two ellipses, whose major axes are equal, have a common focus. Prove that they intersect in two points only.

57. A circle described about the triangle SPS' cuts the minor axes in R on the opposite side to P. Prove that SR varies as the normal PG.

58. If r and R be the radii of the circles inscribed in and about the triangle SPS, prove that $R \,.\, r$ varies as $SP \,.\, S'P$.

59. The circle described upon PG as diameter cuts SP, $S'P$ in K and L. Prove that KL is bisected by PG, and is perpendicular to it.

60. If from S' a line be drawn parallel to SP, it will meet SY in the circumference of a circle.

61. T and t are the points where the tangent at P meets the axes. CP is produced to meet in L the circle described about the triangle TCt; prove that PL is half the chord of the circle of curvature at P in the direction of C, and that $CP \,.\, CL$ is constant.

62. About the triangle PQR an ellipse is described, having its centre at the point where the lines drawn from P, Q, R to the middle points of the opposite sides meet. CP, CQ, CR are produced to meet the ellipse in P', Q', R'. Prove that

the tangents at P', Q', R' form a triangle similar to PQR, and four times as large.

63. Lines from Y and Y' perpendicular to the major axis cut the circles on SP, $S'P$ as diameters in I and J. Prove that IS and JS' when produced, intersect BC in the same point.

64. If from the ends of any diameter chords be drawn to any point in the ellipse, the diameters parallel to these chords will be conjugate.

65. If T be the angle between tangents at the extremities of a focal chord, and O the angle subtended by the chord at the other focus, then

$$2T + O = 2 \text{ right angles.}$$

66. The acute angles which SP, SQ make with the tangents are complementary. Prove that BC^2 is a mean proportional between the areas of the triangles SPS', SQS'. Also, show that the problem is impossible unless $BC < CS$.

67. A series of ellipses have their equal conjugate diameters of the same magnitude. One of these diameters is fixed and common, while the other varies. The tangents drawn from any point in the fixed diameter produced will touch the ellipses in points situated on a circle.

68. If on the longer side of a rectangle as major axis an ellipse be described, passing through the intersection of the diagonals, and lines be drawn from any point of the ellipse exterior to the rectangle to the ends of the remote side, they will divide the major axis into segments, which are in geometric progression.

69. From any point P of an ellipse PQ is drawn at right angles to SP meeting the diameter conjugate to CP in Q. Prove that PQ varies inversely as the perpendicular from P on the major axis.

70. In an ellipse SQ and $S'Q$, drawn at right angles to a pair of conjugate diameters, intersect in Q. Prove that the locus of Q is a concentric ellipse.

CHAPTER III.

THE HYPERBOLA.

45. Def. The *Hyperbola* is the curve traced out by a point which moves in such a manner that its distance from a given fixed point continually bears the same ratio, *greater than unity*, to its distance from a given fixed line. (*See Introduction.*)

Prop. I.

The focus and directrix of a hyperbola being given, to find any number of points on the curve.

Let S be the focus, and MX the directrix.

Draw SX at right angles to the directrix, and divide SX in the point A, so that SA may be to AX in the given fixed ratio, greater than unity; then

A is a point on the curve.

On SX produced take a point A', such that

$$SA' : A'X :: SA : AX;$$

then A' will also be a point on the curve.

On the directrix take *any* point M; and through S and M draw the line $SYMY'$, meeting AY and $A'Y'$, drawn at right angles to AA', in the points Y and Y';

On YY' as diameter describe a circle, and draw PMP parallel to AA', cutting the circle in the points P and P';

P and P' will be points on the hyperbola.



Join PY, PY', SP; then since

$$SY : YM :: SA : AX, \quad (\textit{Euclid}, \text{VI. } 2)$$

$$\text{and } SY' : Y'M :: SA' : A'X, \quad (\textit{Euclid}, \text{VI. } 2)$$

$$\therefore SY : YM :: SY' : Y'M;$$

or, alternately, $SY : SY' :: YM : Y'M$,

and the angle YPY' in a semicircle is a right angle,

$\therefore$ PY bisects the angle SPM,*

$$\therefore SP : PM :: SY : YM,$$

$$:: SA : AX.$$

So we may show that

$$SP' : P'M :: SY' : Y'M,$$

---

* For, if not, make the angle YPm equal to YPS; then

$$SY : Ym :: SP : Pm. \quad (\textit{Euclid}, \text{VI. } 3.)$$

and since, if PY bisect SPm, PY' being at right angles to PY, also bisects the angle between MP and SP produced;

$$\therefore SY' : Y'm :: SP : Pm, \quad (\textit{Euclid}, \text{VI. A.})$$

$$\text{Hence } SY : Ym :: SY' : Y'm,$$

$$\text{or } SY : SY' :: Ym : Y'm,$$

$\therefore$ the points M and m coincide.

$$:: SA : AX,$$

$$\therefore P \text{ and } P' \text{ are points on the curve.}$$

In the same way, by taking other points on the directrix, we may obtain as many more points on the curve as we please.

COR. 1. Since, corresponding to every point P on the curve, there is a point P' situated in precisely the same manner with respect to $A'Y'$ as P is with respect to AY, it is clear that if we make $A'S'$ equal to AS, and $A'X'$ equal to AX, and draw $X'M'$ at right angles to AX', the curve could be equally well described with S' as focus and $M'X'$ as directrix.

The hyperbola is therefore symmetrical, not only with respect to the line AA', but also with respect to the line OC drawn through the middle point of YY' at right angles to and bisecting AA'.

COR. 2. The line OP produced will bisect the angle SPW between SP and $S'P$ produced.

Produce OP and $S'S$ to meet in G. Produce PM to meet $X'M'$ in M', and draw OS' passing through the point M'; then

$$SP : PM :: S'P : PM',$$

$$\text{or, alternately, } SP : S'P :: PM : PM'. \quad (1)$$

$$\text{Again, } SG : PM :: S'G : PM',$$

$$\text{or, alternately, } SG : S'G :: PM : PM', \quad (2)$$

$$\therefore \text{ from (1) and (2)}$$

$$SP : S'P :: SG : S'G,$$

$\therefore PG$ bisects the angle SPW. (*Euclid*, VI. A.)

It will be shown hereafter (*Prop.* IX.) that the normal to the hyperbola at the point P also bisects the angle SPW. Hence the hyperbola and circle have the same tangent at the point P. The hyperbola will consequently touch all the infinite series of circles which can be described in the same manner as the one in the figure, by taking different points on the directrix.

PROP. II.

46. If C be the middle point of AA', then CA is a mean proportional between CS and CX,

$$\text{or } CS \,.\, CX = CA^2. \quad (\textit{See fig. Prop.}\ \text{III.})$$

$$\text{Since } SA' : A'X :: SA : AX.$$

$$\text{Alternately, } SA' : SA :: A'X : AX,$$

$$\therefore\ SA' - SA : SA :: A'X - AX : AX,$$

$$\text{or } AA' : SA :: XX' : AX,$$

$$\therefore\ AA' : XX' :: SA : AX,$$

$$\text{or } CA : CX :: SA : AX. \quad \text{(1.)*}$$

$$\text{Again, } SA' : SA :: A'X : AX.$$

$$\therefore\ SA' + SA : SA :: A'X + AX : AX,$$

$$\text{or } SS' : SA :: AA' : AX.$$

$$\text{Alternately, } SS' : AA' :: SA : AX,$$

$$\text{or } CS : CA :: SA : AX. \quad \text{(2.)}$$

Hence from (1) and (2),

$$CA : CX :: CS : CA,$$

$$\therefore\ CA^2 = CX \,.\, CS.$$

Or CA is a mean proportional between CS and CX.

COR. Since the three lines CS, CA, CX, are proportional, therefore, by the definition of duplicate ratio and *Euclid*, VI. 20 *Cor.*,

$$CS : CX :: CS^2 : CA^2. \quad \text{(3.)}$$

PROP. III.

47. If P be any point on the hyperbola, and S be the focus nearer to P; then

$$S'P - SP = AA'.$$

$$\text{Since } SP : PM :: SA : AX,$$

---

[*] N.B. The results (1), (2), (3), should be remembered, as they will frequently be referred to.



$$\text{and } SA : AX :: AA' : XX', \quad (\textit{Prop.}\ \text{II.})$$

$$\therefore\ SP : PM :: AA' : XX'.$$

$$\text{So } S'P : PM' :: AA' : XX',$$

$$\therefore\ S'P - SP : PM' - PM :: AA' : XX'.$$

But $PM' - PM = MM' = XX'$,

$$\therefore\ S'P - SP = AA'.$$

Cor. By means of this property the hyberbola may be practically described, and the form of the curve determined.

Let a rigid bar $S'Q$ of any length have one end fastened at the focus S', in such a manner that it is capable of turning freely round S' as a centre in the plane of the paper.

At the other end of the bar let a string be fastened of such a length that when stretched along the bar it shall just reach to within a distance equal to AA' from the end S' of the bar.



If the loose end of the string be now fastened to the focus S, and the rod being initially placed in the position $S'S$, be made to revolve round S', while the string is kept constantly stretched by means of the point of a pencil at P, in contact with the bar; since $S'P$ and SP are always increasing by the same amount, viz. the length of the portion of the string that removes itself from the bar, between any two positions of P, the difference between $S'P$ and SP will be constantly the same, and the point P will trace out the hyperbola.

Another perfectly similar branch may be described in the same manner by making the bar revolve round S as centre.

In this case $S'P - SP$ will be equal to AA'.

The curve, therefore, consists of two similar branches, which recede indefinitely both from the line AA', and also from the line BCB' drawn bisecting AA' at right angles. (*See fig. Prop.* IV.)

48. If BC be taken of such a length that

$$BC^2 = CS^2 - CA^2,$$

and CB' be made equal to CB, then AA' and BB' are called respectively the *Transverse* and *Conjugate Axes.*

The line BCB' does not meet the hyperbola, and the reason of its being introduced will be seen further on.

If the conjugate and transverse axes are equal, the hyperbola is said to be *rectangular* or *equilateral.*

The property of the hyperbola, which we have just investigated, viz. that the difference between SP and $S'P$ is constant, is sometimes taken as the definition of the curve. (*See Chapter* II. *Art.* 25.)

Also as in the ellipse, if SL be the semi-latus rectum, it may be proved that

$$SL \,.\, AC = BC^2.$$

PROP. IV.

49. The difference of the distances of any point from the foci of a hyperbola will be greater or less than AA', according as the point is on the concave or convex side of the curve.



(1). Let Q be a point on the concave side of the hyperbola. Join SQ, $S'Q$, and let $S'Q$ meet the curve in P; join SP; then

$$\text{since } S'Q = S'P + PQ,$$
$$\text{and } SQ < SP + PQ,$$



$$\therefore S'Q - SQ > S'P - SP.$$
$$\text{but } S'P - SP = AA',$$
$$\therefore S'Q - SQ > AA'.$$

(2.) Let Q be a point on the convex side of the curve, nearer to S than S'; join SQ, $S'Q$, and let SQ meet the curve in P; join $S'P$; then

$$S'Q < S'P + PQ,$$
$$\text{and } SQ = SP + PQ,$$
$$\therefore S'Q - SQ < S'P - SP,$$
$$\text{but } S'P - SP = AA',$$
$$\therefore S'Q - SQ < AA',$$

so if Q be nearer to S' than S, we can show that

$$SQ - S'Q < AA';$$

Cor. Coversely a point will be on the concave or convex side of the hyperbola, according as the difference of its distances from the foci is greater or less than AA'.

50. Def. If a point P' be taken on the hyperbola near to P (*see fig. Prop.* V.) and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P, is called the *Tangent* to the hyperbola at the point P.

Prop. V.

If the tangent to the hyperbola at any point P meet the directrix in the point Z, and if S be the focus corresponding to the directrix on which Z is situated, then SZ will be at right angles to SP.



Let P' be a point in the curve near to P.

Draw the chord PP', and produce it to meet the directrix in Z; join SZ.

Draw PM, $P'M'$ at right angles to the directrix, and join SP, SP'.

Produce PS to meet the hyperbola in Q; then since the triangles ZMP, $ZM'P'$ are similar,

$$\therefore ZP : ZP' :: MP : M'P',$$
$$:: SP : SP'.$$

$\therefore SZ$ bisects the angle $P'SQ$. (*Euclid*, VI. A.)

Now, when P' is indefinitely near to P, and PP' becomes the tangent at the point P, the angle PSP' becomes indefinitely small, while the angle QSP' approaches two right angles; and therefore the angles ZSP', being half of the angle $P'SQ$, becomes ultimately a right angle.

Hence, when PZ becomes the tangent at the point P, the angle ZSP is a right angle,

or SZ is perpendicular to SP.

Cor. 1. Conversely, if SZ be drawn at right angles to SP, meeting the directrix in Z, and PZ be joined, PZ will be the tangent at P.

Cor. 2. If PZ be produced to meet the other directrix in Z', and $S'Z'$ be joined; then

$S'Z'$ is at right angles to $S'P'$.

Cor. 3. The tangents at the extremities of the latus rectum, or double ordinate through the focus, meet the axis in the point X.

Prop. VI.

The tangent to the hyperbola at any point P makes equal angles with the focal distances SP and $S'P$.



Let the tangent at P meet the directrices in Z and Z'.

Draw PMM' at right angles to the directrices meeting them in M and M' respectively; join SZ, $S'Z'$; then

$$SP : PM :: S'P : PM'.$$

And since the triangles ZMP, $Z'M'P$ are similar,

$$PM : PZ :: PM' : PZ',$$

$$\therefore\ SP : PZ :: S'P : PZ'.\quad (\textit{Ex æquali.})$$

Now in the triangles SPZ, $S'PZ'$ because the sides about the angles SPZ, $S'PZ'$ are proportional, and the angles PSZ, $PS'Z'$ are equal, being right angles, and the angles SZP, $S'Z'P$ are each less than a right angle,

$\therefore$ the triangles SPZ, $S'PZ'$ are similar. (*Euclid*, VI. 7).

$\therefore$ the angle $SPZ = S'PZ'$.

Prop. VII.

The tangents at the extremities of a focal chord intersect in the directrix.

Let PSQ be a focal chord, and let the tangent P meet the directrix in Z. Join SZ; then

the angle ZSP is a right angle, (*Prop.* V.)

And $\therefore$ also the angle ZSQ is a right angle,

$\therefore$ ZQ is the tangent at Q. (*Prop.* V. *Cor.* 1.)

Or the tangents at the extremities of a focal chord intersect in the directrix.

Prop. VIII.

51. If the tangent at P meet the transverse axis in T, and PN be the ordinate of the point P; then

$$CT \cdot CN = CA^2.$$

Draw PMM' at right angles to the directrices meeting them in M and M'. Join SP, $S'P$; then

since PT bisects the angle SPS', (*Prop.* VI.)

$$\therefore\ S'T : ST :: S'P : SP,\quad (\textit{Euclid}, \text{VI. } 3.)$$

$$:: PM' : PM,$$

$$:: X'N : XN.$$



$$\therefore\ S'T - ST : S'T + ST :: X'N - XN : X'N + XN$$
$$\text{or } 2CT : 2CS :: 2CX : 2CN,$$
$$\text{or } CT : CS :: CX : CN.$$
$$\therefore\ CT \cdot CN = CS \cdot CX,$$
$$= CA^2. \quad (\textit{Prop.}\ \text{II.})$$

52. **Def.** The line PG, drawn at right angles to the tangent PT, is called the *Normal* to the hyperbola at the point P.

Prop. IX.

If the normal to the hyperbola at the point P meet the transverse axis in the point G, and PN be the ordinate of the point P, then

$$NG : NC :: BC^2 : AC^2.$$

Draw PMM' at right angles to the directrices, meeting them in M and M', and produce $S'P$ to W; then since

the angle TPG is a right angle,

$\therefore$ the angle WPG = the complement of the angle $S'PT$,

and the angle SPG = the complement of the angle SPT;

but the angle $S'PT =$ the angle SPT,
$\therefore$ the angle $WPG =$ the angle SPG,
$\therefore$ PG bisects the angle SPW,
$\therefore$ $S'G : SG :: S'P : SP$, (*Euclid*, VI. A.)
$:: PM' : PM$,
$:: X'N : XN$,
$\therefore$ $S'G + SG : S'G - SG :: X'N + XN : X'N - XN$;
or $2CG : SS' :: 2CN : XX'$.
Alternately, $2CG : 2CN :: SS' : XX$;
or $CG : CN :: CS : CX$,
$:: CS^2 : CA^2$. (*Prop.* II. *Cor.*)
$\therefore$ $CG - CN : CN :: CS^2 - CA^2 : CA^2$;
or $NG : CN :: BC^2 : AC^2$.

Prop. X.

If PN be the ordinate of any point P on the hyperbola; then

$$PN^2 : AN \,.\, A'N :: BC^2 : AC^2.$$

For $NG : NC :: BC^2 : AC^2$.

And rectangles of the same altitude are to one another as their bases, (*Euclid*, VI. 1.)

$\therefore$ $TN \,.\, NG : TN \,.\, NC :: BC^2 : AC^2$,
or $PN^2 : TN \,.\, NC :: BC^2 : AC^2$
But $TN \,.\, CN = CN^2 - CT \,.\, CN$, (*Euclid*, II. 2)
$= CN^2 - CA^2$, (*Prop.* VIII.)
$= AN \,.\, A'N$, (*Euclid*, II. 6.)
$\therefore$ $PN^2 : AN \,.\, A'N :: BC^2 : AC^2$.

Prop. XI.

If the normal at any point P of an hyperbola meet the transverse axis in G; then

$$SG : SP :: CS : CA.$$

Produce $S'P$ to W; then

since PG bisects the angle SPW, (*Prop.* IX.)

$$SG : S'G :: SP : S'P,$$

$$\therefore SG : S'G - SG :: SP : S'P - SP,$$

but $S'P - SP = AA'$, (*Prop.* III.)

and $S'G - SG = SS'$,

$$\therefore SG : SS' :: SP : AA',$$

$$\text{or } SG : SP :: SS' : AA',$$

$$\text{or } SG : SP :: CS : CA.$$

Cor. Hence also,

$$S'G : S'P :: CS : CA.$$

Prop. XII.

53. If from the foci S and S' of an hyperbola SY and $S'Y'$ are drawn at right angles to the tangent at P, then Y and Y' are on the circumference of the circle described on AA' as diameter, and

$$SY \,.\, S'Y' = BC^2.$$

Join SP, $S'P$, and produce SY to meet $S'P$ in W; join CY; then

since the angle $SPY =$ the angle WPY, (*Prop.* VI.)

and the angle $SYP =$ the angle WYP,

and the side PY is common to the triangles SPY, WPY,

. . the triangle $SPY = WPY$ in all respects,

$$\therefore SP = PW, \text{ and } SY = WY,$$

$$\therefore S'P - SP = S'W,$$

but $S'P - SP = AA'$, (*Prop.* III.)

$$\therefore S'W = AA'.$$

Again, $\therefore SC = CS'$, and $SY = WY$,

$$\therefore SC : CS' :: SY : YW,$$

$$\therefore CY \text{ is parallel to } S'W,$$

$$\therefore CY : SW :: CS :: SS',$$



$$\therefore\ CY = \tfrac{1}{2}S'W = CA;$$

$$\text{so } CY = CA.$$

$\therefore$ Y and Y' are points on the circumference of the circle described upon AA' as diameter.

Next, let SY be produced to meet this circle in Z, and join ZY'; then

since the angle ZYY' is a right angle

$\therefore$ ZY' passes through the centre C,

$\therefore$ the angle SCZ = the angle $S'CY'$,

$$\therefore\ SZ = S'Y',$$

$$\begin{aligned}\therefore\ SY \,.\, S'Y' &= SY \,.\, SZ,\\ &= SA \,.\, SA', \ (\textit{Euclid}, \text{ III. 36 } \textit{Cor.})\\ &= CS^2 - CA^2, \ (\textit{Euclid}, \text{ II. 6.})\\ &= BC^2.\end{aligned}$$

Cor. If CD be drawn parallel to the tangent at P meeting $S'P$ in E; then

since $CEPY$ is a parallelogram,

$$\therefore\ PE = CY = AC.$$

PROP. XIII.

54. To draw a pair of tangents to an hyperbola from an external point O.



Of the foci S and S', let S' be that which is nearer to O.

With centre S and radius equal to AA' describe a circle.

Join OS, OS'; and let SO or SO produced meet the circle in the point I.

Now if O be a point inside the circle MIM' it is evident that OS' is greater than OI; and if O be outside the circle,

$$\text{since } OS - OS' < AA' \text{ or } SI, \quad (\textit{Prop.}\ \text{IV.})$$

$$\therefore OS - OS' < OS - OI,$$

$$\therefore OS' > OI.$$

With centre O and radius OS' describe another circle cutting the former in the points M and M', which it will always do since OS' is greater than OI.

Join SM, SM', and produce them to meet the hyperbola in the points P and P'.

Join OP, OP'; these will be the tangents required.

Join $S'P$, $S'P'$; then

$$\text{since } S'P - SP = AA' = SM,$$
$$\therefore\ S'P = PM.$$
$$\text{And } \therefore\ S'P,\ PO = MP,\ PO, \text{ each to each,}$$
$$\text{and } OS' = OM,$$
$$\therefore \text{ the angle } OPS' = \text{the angle } OPM,$$
$$\therefore\ OP \text{ is the tangent at } P. \quad (\textit{Prop.}\ \text{VI.})$$
$$\text{So } OP' \text{ is the tangent at } P'.$$

The points of contact P and P' will be upon the same or opposite branches of the hyperbola according as SM and SM' require to be produced in the same or in opposite directions with respect to S, in order to intersect the hyperbola.

PROP. XIV.

If from a point O a pair of tangents, OP, OP' be drawn to an hyperbola, then the angles which OP and OP' subtend at either focus will be equal or supplementary according as the points of contact are in the same or opposite branches of the hyperbola.

Let the points P and P' be on opposite branches of the hyperbola.

Join PS, $S'P$; SP', $S'P'$.

Produce PS to M, making PM equal to PS'. Also from $P'S$ cut off a part $P'M'$ equal to $P'S'$.

Join OM, OM'; OS, OS'.

$$\text{Then since } OP,\ PS' = OP,\ PM, \text{ each to each,}$$
$$\text{and the angle } OPS' = \text{the angle } OPM, \ (\textit{Prop.}\ \text{VI.})$$
$$\therefore\ OS' = OM,$$
$$\text{and the angle } OS'P = \text{the angle } OMP.$$



So $OS' = OM'$,

and the angle $OS'P'$ = the angle $OM'P'$,

$\therefore\ OM = OM'$.

Again, $\because\ SM = S'P - SP = AA'$,

and $SM' = SP' - S'P' = AA'$,

$\therefore\ SM = SM'$.

And $\because\ OS, SM = OS, SM'$, each to each,

and $OM = OM'$,

$\therefore$ the angle OSM = the angle OSM',

and the angle OMS = the angle $OM'S$.

But OSM is the supplement of OSP,

and $OM'S$ is the supplement of $OM'P'$,

$\therefore\ OSM'$ is the supplement of OSP,

and OMP the supplement of $OM'P'$.

But $OMP = OS'P$,

and $OM'P' = OS'P'$,

$\therefore\ OS'P$ is the supplement of $OS'P$

Hence the angles which OP and OP' subtend either at S or S' are supplementary.

In a similar manner if P and P' are on the same branch of the hyperbola, the angles subtended either at S or S' may be shown to be equal.

Prop. XV.

55. If the tangent at any point P of an hyperbola meet the conjugate axis in the point t, and Pn be drawn at right angles to CB; then

$$Cn \,.\, Ct = BC^2.$$



Draw PN at right angles to CA; then

$$Ct : CT :: PN : NT,$$
$$\therefore\ Ct : PN :: CT : NT,$$
$$\therefore\ Ct \,.\, Cn : PN^2 :: CT \,.\, CN : CN \,.\, NT;$$
$$\text{or } Ct \,.\, Cn : CT \,.\, CN :: PN^2 : CN \,.\, NT,$$
$$:: BC^2 : AC^2. \quad (\textit{Prop.}\ \text{X.})$$
$$\text{But } CT \,.\, CN = AC^2,$$
$$\therefore\ Ct \,.\, Cn = BC^2.$$

56. The proofs that we have given up to this point of the properties of the hyperbola are closely analogous to the corresponding propositions in the ellipse. The remaining properties of the hyperbola are more conveniently investigated by means of its relation to certain lines, which we shall presently define, called *Asymptotes*, in the same manner as many of the properties of the ellipse were deduced from those of the auxiliary circle.

DEF. The hyperbola described (*see fig. Prop.* XIV.) with C as centre, and BB' as transverse axis, and AA' as conjugate axis, is called the *Conjugate Hyperbola.* Its foci, which will be on the line BCB', will evidently be at the same distance from C as those of the original hyperbola, since

$$CS^2 = CA^2 + CB^2.$$

PROP. XVI.

If through any point R on either of the diagonals of the rectangle formed by drawing tangents to the hyperbola and its conjugate at the vertices, $A, A', B, B',$ two ordinates RPN, RDM, be drawn at right angles to AA' and BB', and meeting either the hyperbola or its conjugate in the points P and D; then

$$RN^2 \backsim PN^2 = BC^2,$$

$$\text{and } RM^2 \backsim DM^2 = AC^2.$$

Let R be a point on the diagonal $O'CO$; then

$$RN^2 : CN^2 :: AO^2 : AC^2,$$

$$:: BC^2 : AC^2,$$

$$\text{and } PN^2 : CN^2 - CA^2 :: BC^2 : AC^2; \text{ (\textit{Prop.} X.)}$$

$$\therefore RN^2 - PN^2 : CA^2 :: BC^2 : AC^2;$$

$$\therefore RN^2 - PN^2 = BC^2.$$

$$\text{Again, } RM^2 : CM^2 :: AC^2 : BC^2,$$

$$\text{and } DM^2 : CM^2 - CB^2 :: AC^2 : BC^2; \text{ (\textit{Prop.} X.)}$$

$$\therefore RM^2 - DM^2 : BC^2 :: AC^2 : BC^2;$$

$$\therefore RM^2 - DM^2 = AC^2.$$



In exactly the same manner, if NR had been produced to meet the conjugate hyperbola in P, and MR had been produced to meet the original hyperbola in D, we should have had,

$$PN^2 - RN^2 = BC^2,$$

$$\text{and } DM^2 - RM^2 = AC^2.$$

COR. If RP be produced to meet the hyperbola in p, and the other asymptote in r; then

$$RN^2 - PN^2 = RP \, . \, Pr \, ; \ (\textit{Euclid}, \text{ II. } 5)$$

$$\therefore \ RP \, . \, Pr = BC^2.$$

Hence as RPN is further removed from A, and the line Pr consequently increases, since the rectangle contained by RP and Pr remains constant, RP must diminish, and by taking R sufficiently far from C, RP may be made less than any assignable magnitude. The line CR, therefore, continually approaches nearer and nearer to the hyperbola, though it never actually reaches it.

On account of this property, CR is called an *Asymptote* to the hyperbola.

So also if P be the point where NR produced meets the conjugate hyperbola, we shall have

$$RP \, . \, Pr = BC^2;$$

and therefore CR is also the asymptote to the conjugate hyperbola.

In the same manner it may be shown that the other diameter oCo' of the rectangle OO' is an asymptote to both hyperbolas.

PROP. XVII.

57. If E be the point where the asymptote meets the directrix; then

$$CE = AC.$$



For by similar triangles,

$$CE : CO :: CX : CA,$$

$$:: CA : CS. \quad (\textit{Prop.}\ \text{II.})$$

But $CO^2 = CA^2 + CB^2 = CS^2$;

$$\therefore CO = CS;$$

$$\therefore CE = AC.$$

Cor. If SE be joined, since

$$CE^2 = CA^2 = CS \,.\, CX,$$

$\therefore$ the angle CES is a right angle. (*Euclid*, VI. 8, *Cor.*)

Prop. XVIII.

If from any point R in one of the asymptotes to an hyperbola ordinates RPN, RDM be drawn to the hyperbola and its conjugate respectively, and PD be joined, PD will be parallel to the other asymptote.

For $RN^2 : RM^2 :: BC^2 : AC^2$;

and $RN^2 - PN^2 : RM^2 - DM^2 :: BC^2 : AC^2$, (*Prop.* XVI).

$$\therefore PN^2 : DM^2 :: BC^2 : AC^4;$$

$$:: RN^2 : RM^2,$$

$$\therefore PN : DM :: RN : RM;$$

$\therefore$ PD is parallel to MN. (*Euclid*, VI. 2.)

Also $CN : CM :: AC : BC$,

$\therefore$ MN is parallel to AB;

and $OA : Ao :: OB : Bo'$,

$\therefore$ AB is parallel to oo'.

Hence PD is parallel to oo'.

Cor. So also if R and D be the points where NR and MR produced meet respectively the conjugate and the original hyperbola, PD will be still parallel to oo'.

PROP. XIX.

58. If through any two points Q and Q' of an hyperbola a line $RQQ'R'$ be drawn in any direction meeting the asymptotes in R and R'; then will

$$RQ = R'Q.$$



Through Q and Q' draw the ordinates UQW, $U'Q'W'$; meeting the asymptotes in U, W, U', W'; then by similar triangles,

$$QR : QU :: Q'R : Q'U',$$

$$\text{and } QR' : QW :: Q'R' : Q'W';$$

∴ compounding

$$QR \,.\, QR' : QU \,.\, QW :: Q'R \,.\, Q'R : Q'U' \,.\, Q'W'.$$

But $QU \,.\, QW = BC^2 = Q'U' \,.\, Q'W$ (*Prop.* XVI. *Cor.*)

$$\therefore QR \,.\, QR' = Q'R \,.\, Q'R';$$

$$\text{but } QR \,.\, QR' = QR \,.\, QQ' + QR \,.\, Q'R',$$
$$\text{and } Q'R \,.\, Q'R' = Q'R' \,.\, QQ' + QR \,.\, Q'R';$$
$$\therefore\ QR \,.\, QQ' = Q'R' \,.\, QQ'.$$
$$\therefore\ QR = Q'R'.$$

COR. 1. If $RQQ'R'$ move parallel to itself until the points Q and Q' coincide, the line RQR' will ultimately assume the position LPl, and will become a tangent to the hyperbola at P.

Hence, since RQ is always equal to $R'Q'$,

$$LP = Pl,$$

or the tangent LPl is bisected at the point of contact P.

COR. 2. If CP be produced to meet RR' in V, then since

$$RV : VR' :: LP : Pl,$$
$$\therefore\ RV = VR';$$
$$\text{and } RQ = Q'R',$$
$$\therefore\ QV = Q'V.$$

Hence if a series of parallel chords be drawn in an hyperbola, their middle points will all be in the line drawn through the centre and the point where the tangent parallel to the chords meets the hyperbola.

DEF. A line PCP' drawn through the centre, and meeting the hyperbola in P and P', is called a *Diameter*.

A diameter consequently bisects all chords drawn parallel to the tangents at its extremities.

PROP. XX.

59. If through any point Q of an hyperbola a line RQr, be drawn in any direction meeting the asymptotes in R and r, and LPl be the tangent drawn parallel to RQr; then

$$RQ \,.\, Qr = PL^2.$$

Through P and Q draw the ordinates EPe, UQW, meeting the asymptote in E, e, U, W; then by similar triangles,



$$QR : QU :: PL : PE,$$
$$Qr : QW :: Pl : Pe$$
$$\therefore QR . Qr : QU . QW :: PL . Pl : PE . Pe$$
but $QU . QW = BC^2 = PE . Pe.$ (*Prop.* XVI. *Cor.*)
$$\therefore QR . Qr = PL . Pl,$$
$$= PL^2.$$ (*Prop.* XIX. *Cor.* 1.)

Cor. If Qq be produced to meet the conjugate hyperbola in Q', q' we may show that

$$Q'R . Q'r = PL^2,$$

and also, as in Proposition XIX., that

$$Q'R = q'r,$$
$$\therefore QQ' = qq'.$$

Hence if a line be drawn in any direction meeting both the hyperbolas, the portions intercepted between the hyperbola and its conjugate will be equal.

PROP. XXI.

60. If from any point P of an hyperbola, PH and PK be drawn parallel to the asymptotes, meeting them in H and K respectively; then $4 \,.\, PH : PK = CS^2$.



Draw the ordinate $RPNr$ meeting the asymptotes in R and r; then by similar triangles,

$$PH : PR :: Co : Oo,$$

$$\text{and } PK : Pr :: CO : Oo,$$

$$\therefore\ PH \,.\, PK : PR \,.\, Pr :: CO^2 : Oo^2,$$

$$:: CS^2 : 4BC^2.$$

$$\text{But } PR \,.\, Pr = BC^2,$$

$$\therefore\ 4 \,.\, PH \,.\, PK = CS^2.$$

PROP. XXII.

If the tangent at any point P of an hyperbola meet the asymptotes in L and l; then the area of the triangle LCl is equal to the rectangle contained by AC and BC.

Draw PH and PK parallel to the asymptotes meeting them in H and K; then

$$\text{since } CL : CH :: Ll : Pl,$$

$$\text{and } Ll = 2\,Pl, \quad (\textit{Prop. } \text{XIX. } \textit{Cor. } 1.)$$

$$\therefore CL = 2\,CH = 2\,PK;$$

$$\text{so } Cl = 2\,CK = 2\,PH,$$

$$\therefore CL \,.\, Cl = 4\,PH \,.\, PK = CS^2, \quad (\textit{Prop. } \text{XXI.})$$

$$= CO \,.\, Co,$$

$$\therefore CL : CO :: Co : Cl,$$

$\therefore$ the triangles LCl, OCo have the angle at C common and the sides about those angles reciprocally proportional;

$$\therefore \text{the triangles } LCl = \text{the triangle } OCo,$$

$$= AC \,.\, AO.$$

$$= AC \,.\, BC.$$

PROP. XXIII.

61. If from any point R in the asymptote of an hyperbola, two ordinates RPN and RDM be drawn to the hyperbola and its conjugate respectively, then the tangents at P and D will be parallel respectively to CD and CP.

Join PD, meeting CR in H; then

$$\text{since } PD \text{ is parallel to } oo', \quad (\textit{Prop. } \text{XVIII.})$$

the tangents at P and D will each meet CR produced in the same point L. (*Prop.* XXII.)

Produce LP and LD to meet the other asymptotes in l and l'; then

$$\text{since } CL \,.\, Cl = CS^2 = CL \,.\, Cl', \quad (\textit{Prop. } \text{XXII.})$$



$$\therefore Cl = Cl',$$

$$\therefore lC : Cl' :: lP : PL,$$

$\therefore$ CP is parallel to the tangent at D.

Also $l'D : DL :: l'C : Cl$,

$\therefore$ CD is parallel to the tangent at P.

The lines CP and CD are called *Conjugate Diameters*, since each of these lines is parallel to the tangent at the extremity of the other.

Prop. XXIV.

If CP and CD be semi-conjugate diameters in the hyperbola; then

$$CP^2 \sim CD^2 = CA^2 \sim CB^2.$$

Draw the ordinates NPR, MDR meeting the asymptote in the point R (*Prop.* XXIII.); then

$$\begin{aligned} CR^2 - CP^2 &= NR^2 - NP^2, \\ &= BC^2, \quad (\textit{Prop.}\ \text{XVI.}) \\ \therefore\ CR^2 &= CP^2 + BC^2; \\ \text{so } CR^2 &= CD^2 + AC^2, \\ \therefore\ CP^2 + BC^2 &= CD^2 + AC^2; \\ \text{or } CP^2 \sim CD^2 &= AC^2 \sim BC^2. \end{aligned}$$

Prop. XXV.

62. The area of any parallelogram formed by drawing tangents to the hyperbola and its conjugate at the extremities P, P', D, D' of a pair of conjugate diameters is equal to the rectangle contained by the axes.



Let $LlL'l'$ be the parallelogram formed by drawing tangents at the extremities P, P', D, D', of any pair of conjugate diameters. The points L, L', l, l', will (*Prop.* XXIII.) be on the asymptotes.

Now the parallelogram $LL' = 4$ parallelogram CL,

$$= 4 \text{ triangle } LCl,$$

$$= 4\, AC \,.\, BC, \quad (\textit{Prop.}\ \text{XXII.})$$

$$= AA' \,.\, BB'.$$

COR. If PF be drawn perpendicular to CD, then

$$PF \,.\, CD = AC \,.\, BC.$$

Also, if the normal PG meet the transverse axis in G, as in the ellipse

$$PF \,.\, PG = BC^2.$$

63. DEF. The line QV drawn from any point Q of the hyperbola parallel to the tangent at any point P, and meeting CP produced in V, is called an *Ordinate* to the diameter CP.

PROP. XXVI.

If QV be an ordinate to the diameter $P'CP$, and CD be conjugate to CP; then

$$QV^2 : PV \,.\, P'V :: CD^2 : CP^2.$$

Produce VQ to meet the asymptotes in R and r; and let the tangent at P meet the asymptotes in L and l; then

$$RV^2 : PL^2 :: CV^2 : CP^2,$$

$$\therefore\ RV^2 - PL^2 : PL^2 :: CV^2 - CP^2 : CP^2.$$

But $RQ \,.\, Qr = PL^2$, (*Prop.* XX.)

$$\therefore\ RV^2 - QV^2 = PL^2,$$

$$\text{or } RV^2 - PL^2 = QV^2.$$

And $CV^2 - CP^2 = PV \,.\, P'V$, (*Euclid*, II. 6.)

$$\therefore\ QV^2 : PL^2 :: PV \,.\, P'V : CP^2.$$

Alternately, $QV^2 : PV \,.\, P'V :: PL^2 : CP^2$.

But since PD is a parallelogram, (*Prop.* XXIII.)

$$\therefore\ PL = CD.$$

Hence $QV^2 : PV \,.\, P'V :: CD^2 : CP^2$.

COR. If VQ be produced to meet the conjugate hyperbola in Q', then

since $Q'R \,.\, Q'r = PL^2$, (*Prop.* XX. *Cor.*)

$$\therefore\ Q'V^2 - RV^2 = PL^2.$$

Hence $Q'V^2 : CV^2 + CP^2 :: CD^2 : CP^2$.

PROP. XXVII.

64. If QV be an ordinate to the diameter PV, and the tangent at Q meet CP in the point T; then

$$CV \,.\, CT = CP^2.$$

Draw the tangent LPl meeting the asymptotes in the points L, l; also let the tangent at Q meet the asymptotes in R, r.



Draw RK, rk parallel to QV meeting CP in K, k.

Now since the triangles RCr, LCl are equal, (*Prop.* XXII.) and have the angle at C common,

$$\therefore\ CR : CL :: Cl : Cr. \quad (\textit{Euclid}, \text{VI. } 15.)$$

$$\text{But } CR : CL :: CK \quad CP,$$

$$\text{and } Cl : Cr :: CP : Ck,$$
$$\therefore CK : CP :: CP : Ck,$$
$$\therefore CK \,.\, Ck = CP^2.$$

Again, produce RK and QV to meet the asymptote Cl in R' and q; then

Since Rr is bisected in Q, (*Prop.* XIX. *Cor.* 1)

$$\therefore R'r \text{ is bisected in } q,$$
$$\text{and } RK = R'K, \text{ (\textit{Prop.} XIX. \textit{Cor.} 2)}$$
$$\therefore Kq \text{ is parallel to } Rr,$$
$$\therefore CT : CK :: Cr : Cq,$$
$$:: Ck : CV,$$
$$\therefore CV \,.\, CT = CK \,.\, Ck$$
$$= CP^2.$$

Cor. 1. Conversely, if QV be an ordinate to PV,

and $CV \,.\, CT = CP^2$,

then QT is the tangent at Q.

Cor. 2. Hence also, if RR' meet the curve in U and U',

and kU, kU' be drawn,

since $CK \,.\, Ck = CP^2$,

kU and kU' are tangents to the hyperbola at U and U'.

Prop. XXVIII.

65. If two chords of a hyperbola intersect one another, the rectangles contained by their segments are proportional to the squares of the diameters parallel to them.

Let QOq be any chord drawn through the point O, and let CD be drawn parallel to it, meeting the conjugate hyperbola in D.

Produce Qq to meet the asymptotes in R and r; and draw the diameter CPV, bisecting both Qq and Rr in V. (*Prop.* XIX. *Cor.* 2.)

Also draw the tangent LPl parallel to Qq, meeting the asymptotes in L and l.



Now since Qq is divided equally in V and unequally in O,

$$\therefore QO \cdot Oq = QV^2 - OV^2; \text{ (\textit{Euclid}, II. 5)}$$

so also $RO \cdot Or = RV^2 - OV^2$, (*Euclid,* II. 5)

$$\therefore RO \cdot Or - QO \cdot Oq = RV^2 - QV^2,$$

$$= RQ \cdot Qr \text{ (\textit{Euclid}, II. 5)}$$

$$= PL^2, \text{ (\textit{Prop.} XX.)}$$

$$\therefore QO \cdot Oq = RO \cdot Or - PL^2.$$

Again, through O and P draw EOe, UPW, at right angles to the axis meeting the asymptotes in E, e, U, W; then

$$RO : OE :: PL : PU,$$

$$\text{and } rO : Oe :: Pl : PW,$$

$$\therefore RO \cdot rO : OE \cdot Oe :: PL^2 : PU \cdot PW;$$

$$\text{but } PU \,.\, PW = BC^2, \text{ (\textit{Prop.} XVI.)}$$
$$\text{and } PL^2 = CD^2, \text{ (\textit{Prop.} XXIII.)}$$
$$\therefore\ RO \,.\, rO : OE \,.\, Oe :: CD^2 : BC^2,$$
$$\text{or } RO \,.\, rO : CD^2 :: OE \,.\, Oe : BC^2,$$
$$\therefore\ RO \,.\, rO - PL^2 : CD^2 :: OE \,.\, Oe - BC^2 : BC^2,$$
$$\text{or } QO \,.\, Oq : CD^2 :: OE \,.\, Oe - BC^2 : BC^2.$$

In the same manner if through O another chord $Q'\,Oq'$ be drawn, and CD' be drawn parallel to it, meeting the conjugate hyperbola in D', we shall have

$$Q'O \,.\, Oq' : CD'^2 :: OE \,.\, Oe - BC^2 : BC^2.$$

Hence $QO \,.\, Oq : Q'O \,.\, Oq' :: CD^2 : CD'^2$.

Cor. The same result may be shown to be true when the point O is outside the hyperbola. Moreover, it is not necessary that the chords should be drawn meeting *one* branch only of the hyperbola or the *same* branch. The proportion still holds good when one or both of the chords meet both branches of the hyperbola, or when the chords are drawn in different branches.

66. Def. If with a point O on the normal at P as centre, and OP as radius, a circle be described touching the hyperbola at P, and cutting it in Q; then when the point Q is made to approach indefinitely near to P, the circle is called the *Circle of Curvature* at the point P.

Prop. XXIX.

If PH be the chord of the circle of curvature at the point P of a hyperbola, which passes through the centre; then

$$PH \,.\, CP = 2\,CD^2.$$

Let PT be the tangent, and PG the normal at the point P.

With centre O, and radius OP, describe a circle cutting the hyperbola in the point Q.

Draw RQW parallel to CP, meeting the circle in W, and TP produced in R.

Also, draw QV parallel to PR, meeting the diameter PP' in V; then since RP touches the circle at P,



$\therefore RQ \,.\, RW = PR^2$, (*Euclid*, III. 36)
or $PV \,.\, RW = QV^2$.
But $QV^2 : PV \,.\, P'V :: CD^2 : CP^2$, (*Prop.* XXVI.)
$\therefore PV \,.\, RW : PV \,.\, P'V :: CD^2 : CP^2$,
or $RW : P'V :: CD^2 : CP^2$.

Now, when the circle becomes the circle of curvature at P, the points R and Q move up to, and coincide with P, and the lines RW and PH become equal, while

$P'V$ becomes equal to PP', or $2CP$.
Hence, $PH : 2CP :: CD^2 : CP^2$,
$\therefore PH \,.\, CP : 2CP^2 :: 2CD^2 : 2CP^2$,
$\therefore PH \,.\, CP = 2CD^2$.

PROP. XXX.

If PU be the diameter of the circle of curvature at the point P of the hyperbola, and PF be drawn at right angles to CD; then

$$PU \,.\, PF = 2CD^2.$$

Since the triangle PHU is similar to the triangle PFC,

$$\therefore PU : PH :: CP : PF,$$

$$\therefore PU \,.\, PF = PH : CP,$$

$$= 2CD^2. \quad (\textit{Prop.}\ \text{XXIX.})$$

Prop. XXXI.

If PI be the chord of the circle of curvature through the focus of the hyperbola; then

$$PI \,.\, AC = 2CD^2.$$

Let $S'P$ meet CD in E; then since the triangles PIU and PEF are similar,

$$\therefore PI : PU :: PF : PE.$$

But $PE = AC$, (*Prop.* XII. *Cor.*)

$$\therefore PI : PU :: PF : AC,$$

$$\therefore PI \,.\, AC = PU \,.\, PF,$$

$$= 2CD^2. \quad (\textit{Prop.}\ \text{XXX.})$$

The point where the circle of curvature intersects the hyperbola may be determined as in the case of the ellipse.

Prop. XXXII.

67. If P be any point on the hyperbola, and CD be conjugate to CP; then

$$SP \,.\, S'P = CD^2.$$

Draw PII' parallel to the asymptote CE meeting the directrices in I and I', and CB' in U.

Let the ordinates, NP, MD meet the asymptote in R, and draw PW perpendicular to the directrix; then by similar riangles,

$$PI : PW :: CE : CX,$$

$$:: CA : CX. \quad (\textit{Prop.}\ \text{XVII.})$$



But $SP : PW :: SA : AX$,

$:: CA : CX$.

$\therefore\ SP = PI$;

so $S'P = PI'$,

$\therefore\ SP \,.\, S'P = PI \,.\, PI'$,

$= UP^2 - UI^2$,

$= CR^2 - CE^2$,

$= CR^2 - CA^2$. (*Prop.* XVII.)

But $CR^2 - CD^2 = RM^2 - DM^2$,

$= CA^2$, (*Prop.* XVI.)

$\therefore\ CR^2 - CA^2 = CD^2$.

Hence $SP \,.\, S'P = CD^2$.

PROBLEMS ON THE HYPERBOLA.

1. THE locus of the centre of a circle touching two given circles is an hyperbola or ellipse.

2. If on the portion of any tangent intercepted between the tangents at the vertices a circle be described, it will pass through the foci.

3. In an hyperbola the tangents at the vertices will meet the asymptotes in the circumference of the circle described on SS' as diameter.

4. If from a point P in an hyperbola PII' be drawn parallel to the transverse axis meeting the asymptotes in I and I'; then $PI \,.\, PI' = AC^2$.

5. If a circle be inscribed in the triangle SPS', the locus of its centre is the tangent at the vertex.

6. If PN be the ordinate of the point P, and NQ a tangent to the circle described on the transverse axis as diameter, and PM be drawn parallel to QC meeting the axis in M, then $MN = BC$.

7. If PN be the ordinate of a point P, and NQ be drawn parallel to AP to meet CP in Q, then AQ is parallel to the tangent at P.

8. If an hyperbola and an ellipse have the same foci, they cut one another at right angles.

9. If the tangent at P intersect the tangents at the vertices in R, r, and the tangent at P' intersect them in R', r', then $AR \,.\, Ar = AR' \,.\, Ar'$.

10. If any two tangents be drawn to an hyperbola, the lines joining the points where they intersect the asymptotes will be parallel.

11. The perpendicular drawn from the focus to the asymptotes of an hyperbola is equal to the semi-conjugate axis.

12. If the asymptotes meet the tangent at the vertex in O, and the directrix in E; then AE is parallel to SO.

13. In a rectangular hyperbola conjugate diameters are equal to one another.

14. In a rectangular hyperbola the normal PG is equal to CP.

15. The lines drawn from any point in a rectangular hyperbola to the extremities of a diameter make equal angles with the asymptotes.

16. Prove that the asymptotes to an hyperbola bisect the lines joining the extremities of conjugate diameters.

17. A line drawn through one of the vertices of an hyperbola and terminated by two lines drawn through the other vertex parallel to the asymptotes will be bisected at the other point where it cuts the hyperbola.

18. P is any point on an hyperbola, and P' a point on the conjugate hyperbola. If CP and CP' be conjugate, prove that

$$S'P' - SP = AC - BC,$$

S and S' being the interior foci.

19. If CP and CD be conjugate, and through C a line be drawn parallel to either focal distance of P, the perpendicular from D upon this line is equal to BC.

20. Given a pair of conjugate diameters, find the principal axes.

21. If Q be a point on the conjugate axis of a rectangular hyperbola, and QP be drawn parallel to the transverse axis meeting the curve in P; then

$$PQ = AQ.$$

22. In a rectangular hyperbola the focal chords drawn parallel to conjugate diameters are equal.

23. If in an equilateral hyperbola CY be drawn at right angles to the tangent at P, and AY be joined, the triangles PCA, CAY are similar.

24. The radius of the circle which touches an hyperbola and its asymptotes, is equal to that part of the latus rectum produced which is intercepted between the curve and the asymptotes.

25. If QQ' by any chord of an hyperbola, and CP the diameter corresponding to it, and QH, PK, $Q'H'$ be drawn parallel to one asymptote meeting the other in H, K and H', then $CH \cdot CH' = CK^2$.

26. If the chord $RPP'R'$ intersect the hyperbola in the points P, P', and the asymptotes in R, R'; and PK be drawn parallel to CR', and $P'K'$ to CR; then $RK = P'K'$, and $R'K' = PK$.

27. If AA' be any diameter of a circle, and PNQ an ordinate to it, then the locus of the intersections of AP, $A'Q$ is a rectangular hyperbola.

28. If two concentric rectangular hyperbolas be described, the axes of one being the asymptotes of the other, they will intersect at right angles.

29. If any chord AP through the vertex be divided in Q, so that $AQ : QP :: AC^2 : BC^2$, and QN be drawn to the foot of the ordinate PN, prove that a straight line drawn at right angles to QN from Q cuts the transverse axis in the same ratio.

30. Prove that the curve which trisects the arcs of all segments of a circle described upon a given base is an hyperbola.

31. If SVs, TVt be two tangents cutting one asymptote in S, T, and the other in s, t, prove that

$$VS : Vs : Vt \; VT.$$

32. If from the exterior focus of an hyperbola a circle be described with radius equal to BC, and tangents be drawn to it from any point in the hyperbola, the line joining the points of contact will touch the circle described on the transverse axis as diameter.

33. Circles are drawn touching the straight line AB in a fixed point C; and from the fixed points A, B tangents are drawn to these circles. The locus of their intersection is an ellipse or hyperbola. Distinguish between the two cases.

34. PP' is a double ordinate in an ellipse. AP, $A'P'$ are produced to meet in Q. Prove that the locus of Q is an hyperbola with the same axes as the ellipse.

35. If the tangent at P intersect the asymptotes in L and l, and PG be the normal at P, then the angle LGl is a right angle.

36. If an ellipse, a parabola, and a hyperbola, have an common tangent, and the same curvature at the vertex, the ellipse will be entirely within the parabola, and the parabola entirely within the hyperbola.

37. The chord $RPP'R'$ of an hyperbola intersects the asymptotes in R and R'. From the point R a tangent RQ is drawn meeting the hyperbola in Q. If PH, QK, $P'H'$ be drawn parallel to one asymptote meeting the other in the points H, K, H'; then $PH + P'H' = 2QK$.

38. If through P, P' on an hyperbola lines be drawn parallel to the asymptotes forming a parallelogram, of which PP is one diagonal; the other diagonal will pass through the centre.

39. If P be the middle point of a line EF which moves so as to cut off a constant area from the corner of a rectangle, its locus is an equilateral hyperbola.

40. PM, PN are drawn parallel to the asymptotes CN, CM, and an ellipse is constructed having CN, CM for semi-conjugate diameters. If CP cut the ellipse in Q, the tangents at Q and P to the ellipse and hyperbola are parallel.

41. If a circle be described through any point P of a given hyperbola and the extremities A, A' of the transverse axis, and NP be produced to meet the circle in Q; prove that Q traces out an hyperbola whose conjugate axis is a third proportional to the conjugate and transverse axes of the original hyperbola.

42. If lines be drawn from any point of a rectangular hyperbola to the extremities of a diameter, the difference between the angles which they make with the diameter will be equal to the angle which this diameter makes with its conjugate.

43. If between a rectangular hyperbola and its asymptotes any number of concentric elliptic quadrants be inscribed, the rectangle contained by their axes will be constant.

44. In the rectangular hyperbola if CP be produced to Q so that $PQ = CP$, and QO be drawn at right angles to CQ to intersect the normal in O, O is the centre of curvature at P.

45. With two conjugate diameters of an ellipse as asymptotes a pair of conjugate hyperbolas are constructed; prove that if one hyperbola touch the ellipse the other will do so likewise; prove also that the diameters drawn through the points of contact are conjugate to each other.

46. If a pair of conjugate diameters of an ellipse when produced be asymptotes to an hyperbola, the points of the hyperbola at which a tangent to the hyperbola will also be a tangent to the ellipse, lie in an ellipse similar to the given one.

47. In the rectangular hyperbola the radius of curvature at P is to the radius of curvature at P' in the triplicate ratio of CP to CP'.

48. OP, OQ are tangents to an ellipse at P and Q, and asymptotes to an hyperbola. Show that a pair of their common chords is parallel to PQ. One of these chords being RS, prove that if PR touches the hyperbola at P, QS touches it at S; also if PS, QR meet in U, OU bisects PQ.

49. The base of the triangle ABC remains fixed, while the vertex C moves in an equilateral hyperbola passing through A and B. If P, Q be the points in which AC, BC meet the circle described on AB as diameter, the intersection of AQ, BP is on the other branch of the hyperbola.

CHAPTER IV.

THE SECTIONS OF THE CONE.

68. Def. If two indefinite straight lines IOI', DOD', intersect one another at a point O, and one of them IOI' remain fixed while the other DOD' revolves round it in such a manner that its inclination to IOI' is the same in all positions, the surface generated by DOD' will be a *Right Cone.*

The line IOI' is called the *Axis*, and the point O the *Vertex* of the *Cone.*

It now remains for us to show (*see Introduction*) that the curve formed by the intersection of this surface with a plane is *in general* one of the three curves whose properties we have been investigating, and to consider under what circumstances it will be the Parabola, Ellipse, or Hyperbola.

If the cutting plane pass through the vertex of the cone as KOK', and intersect the cone again *at all*, it will *in general* cut it in two straight lines as OK, OK' which will represent two positions of the generating line.

The inclination of these lines to each other will depend upon the inclination of the cutting plane to the axis of the cone, and will be greatest when this plane passes through the axis, in which case it will be double the constant angle between the axis and the generating line.

If the cutting plane pass through a generating line dod' and be perpendicular to the plane containing this line and the axis, it will simply touch the cone along this line.



Should the cutting plane not pass through the vertex, and be at right angles to the axis of the cone, the section will evidently be a circle.

In any other case the section will, as we proceed to show, be a Parabola, Ellipse, or Hyperbola.

Whatever be the position of the cutting plane with respect to the cone, we can always suppose a plane drawn through the axis of the cone at right angles to it; and it will be convenient to have this latter plane represented by the plane of the paper as *DOd*. The cutting plane will therefore always be taken at right angles to the plane *DOd* of the parer.

PROP. I.

69. The curve formed by the intersection of the surface of a right cone with a plane (which neither passes through its vertex nor is at right angles to its axis) will be a *Parabola, Ellipse,* or *Hyperbola,* according as the inclination of the cutting plane to the axis of the cone is *equal to*, *greater*, or *less than* the constant angle which the generating line forms with the axis.

Let the plane of the paper represent the plane drawn through the axis IOI' of the cone at right angles to the cutting plane; and let it intersect the surface of the cone in the two generating lines OD, Od.

Let the cutting plane intersect the surface of the cone in the curve PA, and the plane of the paper in the line ANH.

The curve will evidently be symmetrical with respect to this line.

On AH take any point N, and through N draw a plane perpendicular to the axis meeting the surface of the cone in the circle RPr, and the cutting plane in the line PN, which will be at right angles to the plane of the paper and to AN.

Let a sphere be inscribed in the cone touching the cone in the circle EQe and the cutting plane in the point S, and let the plane EQe intersect the cutting plane in the line XM, which will be at right angles to the plane of the paper, and therefore parallel to PN.

Draw PM perpendicular to XM, and join PS, PO, and let PO meet the circle EQe in the point Q.*

Then since PS and PQ are both tangents to the sphere,

$$\therefore PS = PQ.$$

$$\text{But } PQ = RE,$$

$$\therefore PS = RE.$$

---

* N.B. In the figure, to avoid confusion, that part of the section which lies above the plane of the paper is alone represented.



$$\text{But } RE : XN :: AE : AX, \text{ (\textit{Euclid}, VI. 2.)}$$

$$\text{and } AE = AS,$$

$$\therefore RE : XN :: AS : AX,$$

$$\therefore SP : PM :: AS : AX,$$

$\therefore$ the curve PA is either a *Parabola*, *Ellipse*, or *Hyperbola*, whose focus is S and directrix XM.

Again, let AH meet the axis OI in F.

Then the angle AFO will be the inclination of the cutting plane to the axis.

(1) Let the angle AFO = the angle FOd;

then AH is parallel to Od,

$\because$ the angle AXE = the angle OeE = the angle AEX,

$$\therefore AE = AX,$$

$$\therefore AS = AX,$$

$\therefore$ the curve AP will be a *Parabola*.


O
e
E
X
Q
A
M
S
N
F
r
R
P
H
S'
A'
e'
X'
I
E'
D
d
Q'
M'

(2) Let the angle AFO be $> FOd$; then
the complement FXE is $<$ the complement OEe,
$\therefore$ the angle AXE is $< AEX$,
$\therefore AE$ is $< AX$,
or AS is $< AX$,
$\therefore$ the curve AP is an *Ellipse.*

Since the angles HFO, FOd are together less than HFO. OFA, *i.e.* than two right angles, the lines AH and Oe may be produced to meet in A'.

If another sphere be described touching the cone in the circle $E'Q'e'$ and the cutting plane in the point S'; and the line $X'M'$ denote the intersection of the plane $E'Q'e'$ with the cutting plane, and PM' be drawn at right angles to this line, it can easily be shown that

$$S'P : PM' :: S'A' : A'X'.$$

Hence S' and $X'M'$ represent respectively the other focus and directrix of the ellipse.

Also if BC be the semi-axis minor, and through the centre C a line UCU' be drawn parallel to Ee meeting OD, Od in U and U', then it is evident that

$$BC^2 = CU \, . \, CU'.$$

(3) Let the angle AFO be $< FOd$; then
the angle AXE is $>$ the angle AEX,
$\therefore AE$ is $> AX$,
$\therefore AS$ is $> AX$,
$\therefore$ the curve PA is an *Hyperbola.*

Since the angles AFO, FOd' are less than the two FOd, FOd', *i.e.* than two right angles, the lines FA and dO may be produced to meet in A'.

In this case the cutting plane will intersect the other half of the cone, and if any point P' be taken on this part of the curve, and $P'M$ be drawn at right angles to XM, it can be shown as before that

$$SP' : P'M :: S'A :: AX.$$


F'
d'
D'
S
A'
F'
X'
e'
Q'
M'
C
O
e
E
X
Q
A
S
M
r
L
I
N
R
d
K
P
D
F

The intersection of the cutting plane therefore with this portion of the cone will be the other branch of the hyperbola.

Also if another sphere be described touching the upper portion of the cone in $E'Q'e'$, and the cutting plane in S', and the line $X'M'$ denote the intersection of the plane $E'Q'e'$ with the cutting plane, and $P'M'$ be drawn at right angles to this line, it can be easily shown that

$$S'M' : P'M' :: S'A' : A'X'.$$

Hence, S' and $X'M'$ will represent respectively the other focus and directrix of the hyperbola.

Cor. 1. In this last case, *i.e.* when the section is an hyperbola, if a plane OKL be drawn through the vertex of the cone parallel to the cutting plane, meeting the plane of the paper in the straight line OL, and the surface of the cone in the generating line OK; then

$$\begin{aligned} OL : OK &:: OL : OR, \\ &:: AN : AR, \\ &:: AX : AE, \text{ (\textit{Euclid}, VI. 2)} \\ &:: AX : AS, \\ &:: CA : CS, \text{ (\textit{Chap.} III. \textit{Prop.} II.)} \end{aligned}$$

where C is the middle point of AA', and therefore the centre of the hyperbola.

$\therefore$ KOL is half the angle between the asymptotes. (*Chap.* III. *Prop.* XVI.)

Again, if BC be the conjugate semi-axis, and $CU'U$ be drawn parallel to Rr meeting OD', Od' in U and U', then

$$\begin{aligned} \text{since } CU : AC &:: RL : OL, \\ \text{and } CU' : A'C &:: rL : OL, \\ \therefore CU.CU' : AC^2 &:: RL.rL : OL^2, \\ &:: KL^2 : OL^2; \\ \text{but } BC^2 : AC^2 &:: KL^2 : OL^2, \\ \therefore BC^2 &= CU.CU'. \end{aligned}$$

Cor. 2. If the cutting plane is parallel to the axis OL and OI coincide.



In this case half the angle between the asymptotes of the hyperbolic section is equal to the constant angle DOI, and we can at once see that OC is the semi-conjugate axis.

This affords a convenient method of obtaining a pair of conjugate hyperbolas.

Draw Oi at right angles to OI in the plane of the paper, and let another cone be formed by supposing OD to revolve round Oi in such a manner that the angle DOi is the same in all positions, and equal to the complement of DOI.

Then if through any point A on the common generating line OD we draw two planes at right angles to the plane of the paper, and parallel respectively to OI and Oi, they will cut the cones in two hyperbolas, whose semi-transverse axes will be respectively AC, OC, and whose semi-conjugate axes will be respectively OC, AC, and which therefore will be *conjugate* to each other.

70. As long as the cutting plane remains parallel to itself, it is evident that the ratio of AE to AX, and therefore of AS to AX will be altered. Hence the sections made by planes inclined at the same angle to the axis of the cone will have the same eccentricity.*

71. Through any point Q on the circle EQe let a plane be drawn parallel to ANP, intersecting the plane of the paper in the straight line WLN', the cone in the curve WQP', and the planes of the circular sections EQe, RPr in the ordinates QL, $P'N'$.



Then it is manifest that the curve WQP' will touch the circle wQw', formed by the intersection of the cutting plane with the sphere at the extremities of the ordinate QL produced.

Join OP' meeting EQe in Q'; then

$$P'Q' : N'L :: RE : NX,$$
$$:: SA : AX.$$

---

* The ratio of $SA : AX$, or of $CS : CA$, is called the eccentricity.

But $P'Q'$ is equal to the tangent drawn from P' to the circle wQw', and $N'L$ is equal to the perpendicular from P' on the common ordinate of the circle wQw' and the section WQP'.

Hence we have the following important property,—

If a circle touch a conic section in two points at the extremities of an ordinate, the ratio which the tangent drawn to the circle from any point on the curve bears to the perpendicular from the same point on the common chord is equal to the eccentricity of the conic section.

If the two points in which the circle touches the conic coincide, the circle becomes the circle of curvature at the vertex; and therefore the ratio which the tangent drawn from any point of a conic section to the circle of curvature at the vertex bears to the abscissa of the point, is constant and equal to the eccentricity of the curve.

PROBLEMS ON THE SECTIONS OF THE CONE.

1. THE foci of all parabolic sections which can be cut from a given right cone lie upon the surface of another cone.

2. The foci of all elliptical sections of a given right cone, in which the ratio of CA to CS is the same, will lie on two other cones.

3. The extremities of the minor axes of the elliptical sections of a right cone made by parallel planes lie on two generating lines.

4. The latus rectum of a parabola cut from a given cone varies as the distance between the vertices of the cone and the parabola.

5. Under what conditions is it possible to cut an equilateral hyperbola from a given right cone?

6. Two cones whose vertical angles are supplementary are joined as in *Art.* 69, *Cor.* 2. Prove that the latera recta of the curves of section of the cones, whose axes are respectively OI and Oi, made by planes parallel or perpendicular to the plane of the axes, are in the duplicate ratio of Oi and OI.

ADDITIONAL PROBLEMS.

1. Show that the part of the directrix of a parabola, intercepted between the perpendiculars on it from the extremities of any focal chord, subtends a right angle at the focus.

2. The locus of the foci of all parabolas touching the three sides of a triangle is a circle. Prove this, and give a geometrical construction for finding the centre.

3. A system of parabolas which always touch two given straight lines have their axes parallel; show that the locus of the foci is a straight line.

4. Prove that the locus at the foot of the perpendicular from the focus of a parabola on the normal is a parabola.

5. If S be the focus of a parabola, which touches the sides AB, AC of the triangle ABC at the points B, C, and O the centre of the circle described about the triangle; prove that the angle OSA is a right angle.

6. From the focus of a parabola a straight line is drawn parallel to the tangent at any point P, meeting the diameter through P in V'; show that the tangent drawn from P to any circle passing through V' is equal to one-half of the ordinate QV, V being the second point in which the circle cuts the diameter through P.

7. PSp is a focal chord, and upon PS and pS as diameters, circles are described; prove that the length of either of their common tangents is a mean proportional between AS and Pp.

8. If AQ be a chord of a parabola through the vertex A, and QR be drawn perpendicular to AQ to meet the axis in R; prove that AR will be equal to the chord through the focus parallel to AQ.

9. The locus of the vertices of all parabolas, which have a common focus and a common tangent, is a circle.

10. Two parabolas have a common axis and vertex, and their concavities turned in opposite directions; the latus rectum of one is eight times that of the other; prove that the portion of a tangent to the former, intercepted between the common tangent and axis, is bisected by the latter.

11. B is a point on a radius OA of a circle, whose centre is O. On OA produced a point C is taken, such that $OB \cdot OC = OA^2$. If P be any point on the circumference of this circle, R the middle point of BP, and Q the point of intersection of AR, CP; prove that the locus of Q is a circle.

12. If from the middle point of a focal chord of a parabola two straight lines be drawn, one perpendicular to the chord meeting the axis in G, and the other perpendicular to the axis meeting it in N; show that NG is constant.

13. A circle is drawn touching the axis of a parabola, the focal distance of a point P, and the diameter through P. Show that the locus of its centre is a parabola with vertex S, and latus rectum equal to AS.

14. If from the point of intersection of the directrix and axis of a parabola a chord XPQ be drawn, cutting the parabola in P, Q; show that the rectangle contained by the ordinates of PQ is equal to the square of one-half the latus rectum.

15. Find the locus of the centre of a circle which touches a given circle and a given straight line.

16. Given one point of contact of a parabola with three

tangents given in position, find the two other points of contact.

17. The triangle ABC circumscribes a parabola whose focus is S. Through A, B, C, lines are drawn perpendicular respectively to SA, SB, SC. Show that these lines pass through one point.

18. From the focus a line is drawn parallel to the tangent at P, meeting the parabola at Q. QN is an ordinate, and the tangents at P and Q meet the axis in T and T'. Prove that $SN^2 = 4AT \,.\, AT''$, and that if the diameter at P meet SQ in E, the locus of E is a parabola, whose latus rectum is half that of the given parabola.

19. P is any point in a parabola; through S a line is drawn at right angles to the axis, meeting the chord AP or AP produced in R. Prove that $SK \,.\, SR = 2AS \,.\, SY$, where SY is the perpendicular on the tangent, and SK on the normal.

20. From the focus S of a parabola SK is drawn, making a given angle with the tangent at P. Show that the locus of K is that tangent to the parabola which makes with the axis an angle equal to the given angle.

21. PSQ is a focal chord of a parabola, AP' a parallel chord meeting the latus rectum in Q'; prove that $AP' \,.\, AQ = SP \,.\, SQ$.

22. The circle of curvature at any point of a parabola whose abscissa is AN cuts the axis in U and U'. Prove that $AU \,.\, AU' = 3AN^2$.

23. AB is a diameter of a circle. From any point Q in the circumference a tangent QP is drawn, and from P a perpendicular PN is let fall upon AB. Show that if P be always taken so that QP is equal to AN, the locus of P will be a parabola.

24. If a tangent be drawn from any point of a parabola to the circle of curvature at the vertex, the length of the tangent

will be equal to the abscissa of the point measured along the axis.

25. To two parabolas which have a common focus and axis two tangents are drawn at right angles; the locus of their intersection is a straight line parallel to the directrices.

26. If any three tangents be drawn to a parabola, the circle described about the triangle so formed will pass through the focus, and the perpendiculars from the angles on the opposite sides intersect in the directrix.

27. A parabola touches one side of a triangle in its middle point, and the other two sides produced. Prove that the perpendiculars drawn from the angles of the triangle upon any tangent to the parabola are in harmonical progression.

28. Two equal parabolas have the same axis and vertex, but are turned in opposite directions; chords of one parabola are tangents to the other. Show that the locus of the middle point of the chords is a parabola whose latus rectum is one-third of that of the given parabola.

29. Two equal parabolas have the same focus, and their axes are at [right angle to each other, and a normal to one of them is perpendicular to a normal of the other; prove that the locus of the intersection of such lines is a parabola.

30. Show that in every ellipse there are two equal conjugate diameters, coinciding in direction with the diagonals of the rectangle, which touches the ellipse at the extremities of the axes.

31. If a circle be described through the two foci of an ellipse, cutting the ellipse, show that the angle between the tangents to this circle, and to the ellipse at either point of intersection, is equal to the inclination of the normal to the ellipse to the axis minor.

32. The points in which the tangents at the extremities of the transverse axis of an ellipse are cut by the tangent at any

point of the curve, are joined one with each focus; prove that the point of intersection of the joining lines lies in the normal at the point.

33. The external angle between any two tangents to an ellipse is equal to the semi-sum of the angles which the chord joining the points of contact subtends at the foci.

34. The tangent to an ellipse at any point P is cut by any two conjugate diameters in T, t, and the points T, t, are joined with the foci S, S' respectively; prove that the triangles SPT, $S'Pt$ are similar to each other.

35. P is any point on a fixed circle, the centre of which is O; E is a fixed point without the circle; an ellipse is described with centre O and area constant so as always to touch EP at P. Find the locus of the extremities of the diameter conjugate to OP.

36. The normal at any point P of an ellipse cuts the axes in G, g; prove that if any circle be described passing through G, g, the tangent to it from P is equal to CD.

37. Given a focus, a tangent, and the eccentricity of a conic section; prove that the locus of the centre is a circle.

38. A straight line is drawn through a given point C within a circle to cut it in P, P'. If p is taken in it such that $Cp^2 = CP \, . \, CP'$, find the locus of p.

39. In the ellipse $PY \, . \, PY' : PN^2 :: CS^2 : BC^2$ and $SY \, . \, CD = SP \, . \, BC$.

40. Show that if the distance between the foci of the ellipse be greater than the length of its axis minor, there will be four positions of the tangent, for which the area of the triangle, included between it and the straight lines drawn from the centre of the curve to the feet of the perpendiculars from the foci on the tangent, will be the greatest possible.

41. Two conjugate diameters of an ellipse are cut by the tangent at any point P in M, N; prove that the area of the triangle CPM varies inversely as that of the triangle CPN.

42. Circles are described on SY, $S'Y'$ as diameters, cutting SP, $S'P$ respectively in Q, Q'. Prove that $SQ \cdot S'P = SP \cdot S'Q' = BC^2$.

43. PSP', pSp' are any two focal chords of a conic section, P and p being on the same side of the axis; prove that Pp, $P'p'$ meet on the directrix.

44. Prove that an ellipse can be inscribed in any parallelogram so as to touch the middle points of the four sides; and show that this ellipse is the greatest of all inscribed ellipses.

45. If from any point on the exterior of two concentric, similar, and similarly placed ellipses, two tangents be drawn to the interior ellipse which also intersect the exterior; show that the distance between the points of intersection will be double of that between the points of contact.

46. The tangent at any point P in an ellipse, of which S and H are the foci, meets the axis major in T, and TQR bisects HP in Q, and meets SP in R; prove that PR is one-fourth of the chord of curvature at P through S.

47. Prove that the distance between the two points on the circumference of an ellipse at which a given chord, not passing through the centre, subtends the greatest and least angles, is equal to the diameter which bisects that chord.

48. From any point on the auxiliary circle chords are drawn through the foci of an ellipse, and straight lines join the extremities of the chords with the extremity of the diameter passing through the point; prove that these lines will touch the ellipse.

49. A quadrilateral circumscribes an ellipse. Prove that either pair of opposite sides subtends supplementary angles at either focus.

50. Two tangents to an ellipse intersect at right angles; show that the straight line joining their point of intersection with the point of intersection of the normals at the points of contact passes through the centre.

51. P, Q are points in two confocal ellipses, at which the line joining the common foci subtends equal angles; prove that the tangents at P, Q are inclined to an angle which is equal to the angle subtended by PQ at either focus.

52. Tangents to an ellipse are drawn from any point in a circle through the foci; prove that the lines bisecting the angle between the tangents all pass through a fixed point.

53. If the ordinate at P meet the auxiliary circle in Q, and CQ meet the ellipse in R, then CR is equal to the perpendicular on the tangent at P from C.

54. If P be a point such that SP, $S'P$ are perpendicular; prove that $CD^2 = 2 \,.\, BC^2$.

55. If circles be described to the triangle SPS' opposite to the angles S and S'; prove that the rectangle contained by their radii is equal to BC^2.

56. The circle of curvature at any point P of an ellipse meets the focal distances in R, R'; SU is a tangent to the circle.

Prove that $SU^2 : SP^2 :: 2 \,.\, SP - 3 \,.\, AC : AC$,

and if RR' passes through the centre of the circle of curvature, $CP = CS$. Determine the limits of possibility in both cases.

57. A straight line is drawn from the centre of an ellipse meeting the ellipse in P, the circle on the major axis in Q, and the tangent at the vertex in T. Prove that as CT approaches and ultimately coincides with the semi-major axis, PT and QT are ultimately in the duplicate ratio of the axes.

58. A straight line is drawn through the focus S of an ellipse meeting the two tangents at right angles to it in Y and Z, the diameter parallel to these tangents in L, and the directrix in M; prove that

$$SL : SY :: SZ : SM.$$

59. If any equilateral triangle PQR be described in the auxiliary circle of an ellipse, and the ordinates to P, Q, R meet the ellipse in P', Q', R'; the circles of curvature at P', Q', R', meet in one point lying on the ellipse.

60. From a point T two tangents TP, TQ are drawn to an ellipse. Show that a circle with T as centre can be described so as to touch SP, $S'P$, SQ, $S'Q$.

61. If the normal at P meet the axis minor in g, and if the tangent at P meet the tangent at the vertex A in V; show that

$$Sg : SC :: PV : VA.$$

62. If a circle passing through Y and Y' touch the major axis in Q, and that diameter of the circle which passes through Q meet the tangent in P; show that $PR = BC$. (*See fig. Prop.* XV.)

63. If PG the normal at P cut the major axis in G, and if DR, PN be the ordinates of D and P, prove that the triangles PGN, DRC are similar; and thence deduce that PG bears a constant ratio to CD.

64. The tangent at a point P of an ellipse meets the tangents at the vertices in V, V'. On VV' as diameter, a circle is described which intersects the ellipse in Q, Q'; show that the ordinate of Q is to the ordinate of P as BC to $BC + CD$; where CD is conjugate to CP.

65. PCP' is any diameter of an ellipse; the tangents at any two points E and E' intersect in F; PE', $P'E$ intersect in G. Show that FG is parallel to the diameter conjugate to PCP'.

66. If P be any point on an ellipse, and with P as centre and the semi-axis minor as radius a circle be described; prove that if PG be the normal, a circle described on CG as diameter will cut the first circle at right angles.

67. ABC is an isosceles triangle having $AB = AC$. BD, BE drawn on opposite sides of BC, and equally inclined to it, meet AC in D and E.

If an ellipse be described about BCD having its minor axis parallel to BC; then AB will be a tangent to the ellipse.

68. If AQ be drawn from one of the vertices of an ellipse perpendicular to the tangent at any point P; prove that the locus of the point of intersection of PS and QA produced will be a circle.

69. If Y, Y' be the feet of the perpendiculars from the foci of an ellipse on the tangent at P; prove that the circle circumscribed about the triangle YNY' will pass through C.

70. Prove that the angle between the tangents to the auxiliary circle at Y, Y', is the supplement of the angle SPS'.

71. P is any point on an ellipse; PM, PN perpendicular to the axes meet respectively, when produced, the circles described on the axes as diameter in the points Q, Q'. Show that QQ' passes through the centre.

72. Assuming that the greatest triangle which can be inscribed in a circle is equilateral, prove by the method of projection, that the greatest triangle which can be inscribed in an ellipse has one of its sides bisected by a diameter of the ellipse, and the others cut in points of bisection by the conjugate diameter.

73. PQ is a chord of an ellipse, normal at P, LCL' the diameter bisecting it. Show that PQ bisects the angle LPL', and that $LP + L'P$ is constant.

74. A tangent to an ellipse at a point P intersects a fixed tangent in T; if through the focus a line be drawn perpendicular to ST meeting the tangent to P in Q; show that the locus of Q is a straight line touching the ellipse.

75. In an ellipse if a line be drawn through the focus making a constant angle with the tangent; prove that the locus of the point of intersection with the tangent is a circle.

76. Any chord PP' of an ellipse is produced to a point Q, such that $P'Q$ is equal to half the diameter parallel to PP', and $QR'R$ is drawn through the centre to meet the ellipse in R, R'; show that the area PCR is three times the area PCR.

77. In an ellipse, L is the extremity of the latus rectum, and CD conjugate to CL. If a circle be described with centre C and passing through B, and a line be drawn through D parallel to the major axis, the portion of this line which lies within the circle will be equal to the latus rectum.

78. If P be any point in an ellipse, and K the point in which a normal at P intersects a line at right angles to it through S', E the point of intersection of SP, and the diameter conjugate to CP, and if EK and CK be joined, each of the figures $SCKE$, $S'CEK$ will be a parallelogram.

79. If T be a point on the axis AA' produced, and PN the ordinate of the point where the tangent from T touches the ellipse; prove that

$$AN \,.\, A'N : AT \,.\, A'T :: CN : CT.$$

80. Given in an ellipse a focus and two tangents; prove that the locus of the other focus is a straight line.

81. A focus, a tangent, and the axis major being given, prove that the locus of the other focus is a circle.

82. A focus, a tangent, and the axis minor being given, prove that the locus of the other focus is a straight line.

83. An ellipse touches a fixed ellipse and has a common focus with it; if the major axis be fixed, the locus of the other focus is a circle; if the minor axis be fixed, the locus is an ellipse.

84. An ellipse and a parabola have a common focus. Prove that the ellipse either intersects the parabola in two points, and has two common tangents with it, or else does not cut it.

85. If in the ellipse a focus, a point, and the axis minor be given, the locus of the other focus is a parabola.

86. If at the extremities P, Q of any two diameters CP, CQ of an ellipse, two tangents Pp, Qp be drawn cutting each other in T, and the diameter produced in p and q, then the areas of the triangles TQp, TPq are equal.

87. If a straight line CN be drawn from the centre to bisect that chord of the circle of curvature at any point P of an ellipse, which is common to the ellipse and circle, and if it be produced to cut the ellipse in Q, and the tangent in T; prove that $CP = CQ$, and that each is a mean proportional between CN and CT.

88. An ellipse is described so as to touch the three sides of a triangle; prove that if one of its foci move along the circumference of a circle passing through two of the angular points of the triangle, the other will move along the circumference of another circle, passing through the same two angular points. Prove also that if one of these circles pass through the centre of the circle inscribed in the triangle, the two circles will coincide.

89. A triangle is described about an ellipse, so that the extremities of one of its sides lie in an ellipse, confocal with the given one; prove that the line bisecting the opposite angle passes through the pole of that side with respect to the outer ellipse.

90. Prove the following construction for a pair of tangents from any external point T to an ellipse of which the centre is C. Join CT; let $TPCP'T$, a similar and similarly situated ellipse, be drawn, of which CT is a diameter, and P, P' its points of intersection with the given ellipse; TP, TP' will be tangents to the given ellipse.

91. The locus of the foci of all ellipses inscribed in the same parallelogram is a rectangular hyperbola. Prove this, and give a geometrical construction for finding the asymptotes.

92. AC is a fixed diameter of a circle, $ABCD$ a quadrilateral figure inscribed in the circle; prove that if the angles BAC, DAC be complementary, the locus of the intersection of BA, CD will be an hyperbola.

93. Prove that a circle can be described so as to touch the four straight lines drawn from the foci of an hyperbola to any two points on the same branch of the curve.

94. Any three diameters of an ellipse LL', MM', NN', being taken, a circumscribing parallelogram $RTUV$ touches the ellipse at L, L' M, M'. Show that a conic section can be described through the points R, T, U, V, N, N', which will be an hyperbola whose asymptotes are the lines forming in the ellipse the diameters conjugate to NN' and to the other common chord of the ellipse and hyperbola.

95. On opposite angles of any chord of a rectangular hyperbola are described equal segments of circles; show that the four points in which the circles to which these segments belong again meet the hyperbola, are the angular points of a parallelogram.

96. A triangle is inscribed in a rectangular hyperbola: prove that the circle described through the middle points of the sides of the triangle passes through the centre of the hyperbola.

97. ACB is an isosceles triangle; AB the base, and D any point in CB or CB produced: if BZ be drawn parallel to AD, meeting CA or CA produced in Z, prove that the middle point of DZ will be in an hyperbola whose asymptotes are CA, CB.

98. An ellipse and hyperbola are described so that the foci of each are at the extremities of the transverse axis of the other; prove that the tangents at their points of intersection meet the conjugate axis in points equidistant from the centre.

99. In a rectangular hyperbola, PK, PL are drawn at right angles to $A'P$, AP respectively to meet the transverse axis in K and L; prove that PK is equal to AP and KL to AA', and the normal at P bisects KL.

100. In a rectangular hyperbola PC is a fixed diameter, Q any point on the curve; show that the angles QPO, QOP differ by a constant angle.

101. If the tangent at any point P of an hyperbola cut an asymptote in T, and if SP cut the same asymptote in Q, when $SQ = QT$.

102. If a given point be the focus of any hyperbola, passing through a given point and touching a given straight line, prove that the locus of the other focus is an arc of a fixed hyperbola.

103. At any P of an hyperbola a tangent is drawn, and PQ is taken on it in a constant ratio to CD; prove that the locus of Q is an hyperbola.

104. In an hyperbola, supposing the two asymptotes and one point of the curve be given in position, show how to construct the curve; and find the position of the foci.

105. If A, D be two fixed points, and the angle PAD always exceed PDA by a given angle; find the locus of P, and the position of the transverse axis and asymptote.

106. From the middle point D of the base AB of the triangle ABC a straight line EDE' is drawn, making a given angle with AB, and the points E, E' are taken so that $ED = E'D = \frac{1}{2} AB$. If CA, CB take all possible positions consistent with the condition that the difference of the angles CAB, CBA is equal to EDA; prove that the point C will trace out a rectangular hyperbola of which AB, $E'E$ are conjugate diameters.

107. In the rectangular hyperbola, prove that the triangle, formed by the tangent at any point and its intercepts on the axes, is similar to the triangle formed by the straight line joining that point with the centre, and the abscissa and semi-ordinate of the point.

108. Tangents are drawn to an hyperbola, and the portion of each tangent intercepted by the asymptotes is divided in a constant ratio; prove that the locus of the point of section is an hyperbola.

109. Show that the point of trisection of a series of conterminous circular arcs lie on branches of two hyperbolas, and determine the distance between their centres.

110. From a point R on one asymptote RE is drawn touching the hyperbola in E, and ET, EV are drawn through E, parallel to the asymptotes, cutting a diameter in T and V; RV is joined, cutting the hyperbola in P, p: show that TP, Tp touch the hyperbola.

111. Given in the ellipse a focus and two points, the locus of the other focus is an hyperbola.

112. If a rectangular hyperbola passes through three given points, the locus of its centre is a circle, which passes through the middle points of the lines joining the three given points.

113. If the tangent at P meet one asymptote in T, and a line TQ be drawn parallel to the other asymptote to meet the curve in Q; prove that if PQ be joined and produced both ways to meet the asymptotes in R and R', RR' will be trisected at the points P and Q.

114. If two concentric rectangular hyperbolas have a common tangent, the lines joining their points of intersection to their respective points of contact with the common tangent will subtend equal angles at their common centres.

115. If TP, TQ be two tangents drawn from any point T to touch a conic in P and Q, and if S and S' be the foci, then

$$ST^2 : S'T^2 :: SP \,.\, SQ : S'P \,.\, S'Q.$$

116. The circle of curvature at the vertex of a conic meets the axis again in D, and a tangent is drawn to the circle at D: if two tangents be drawn to the circle from any point in the conic they will intercept between them a constant length of the former tangent.

117. If the lines which bisect the angles between pairs of tangents to an ellipse be parallel to a fixed straight line, prove that the locus of the points of intersection of the tangents will be a rectangular hyperbola.

118. An hyperbola, of given eccentricity, always passes through two given points; if one of its asymptotes always pass through a third given point in the same straight line with these, prove that the locus of the centre of the hyperbola will be a circle.

119. A, P and B, Q are points taken respectively in two parallel straight lines, A and B being fixed, and P, Q variable. Prove that if the rectangle $APBQ$ be constant, the line FQ will always touch a fixed ellipse or a fixed hyperbola, according as P and Q are on the same or opposite sides of AB.

120. If two plane sections of a right cone be taken, having the same directrix, the foci corresponding to that directrix lie on a straight line which passes through the vertex.

121. Give a geometrical construction by which a cone may be cut so that the section may be an ellipse of given eccentricity.

122. Given a right cone and a point within it, there are but two sections which have this point for focus; and the planes of these sections make equal angles with the straight line joining the given point and the vertex of the cone.

123. If the curve formed by the intersection of any plane with a cone be projected upon a plane perpendicular to the axis, prove that the curve of projection will be a conic section, having its focus at the point in which the axis meets the plane of projection.

124. If F be the point where the major axis of an elliptic section meets the axis of the cone, and C be the centre of the section; prove that

$$CF : CS :: AA' : AO + A'O,$$

O being the vertex of the cone.

THE END.

R. CLAY, SONS, AND TAYLOR, PRINTERS, BREAD STREET HILL.

SEPTEMBER, 1869.

A CATALOGUE of EDUCATIONAL BOOKS, with a Short Account of their Character and Aim,

Published by

MACMILLAN AND CO.

Bedford Street, Covent Garden, London.

CLASSICAL.

ÆSCHYLI EUMENIDES. The Greek Text, with English Notes and English Verse, Translation, and an Introduction. By BERNARD DRAKE, M.A., late Fellow of King's College, Cambridge. 8vo. 3*s.* 6*d.*

The Greek text adopted in this Edition is based upon that of Wellauer, which may be said, in general terms, to represent that of the best manuscripts. But in correcting the Text, and in the Notes, advantage has been taken of the suggestions of Hermann, Paley, Linwood, and other commentators. In the Translation, the simple character of the Æschylean dialogues has generally enabled the author to render them without any material deviation from the construction and idioms of the original Greek.

ARISTOTLE ON FALLACIES; OR, THE SOPHISTICI ELENCHI. With a Translation and Notes by EDWARD POSTE, M.A., Fellow of Oriel College, Oxford. 8vo. 8*s.* 6*d.*

Besides the doctrine of Fallacies, Aristotle offers, either in this treatise or in other passages quoted in the commentary, various glances over the world of science and opinion, various suggestions or problems which are still agitated, and a vivid picture of the ancient system of dialectics, which it is hoped may be found both interesting and instructive.

Aristotle. — AN INTRODUCTION TO ARISTOTLE'S RHETORIC. With Analysis, Notes, and Appendices. By E. M. COPE, Senior Fellow and Tutor of Trinity College, Cambridge. 8vo. 14*s.*

This work is introductory to an edition of the Greek Text of Aristotle's Rhetoric, which is in course of preparation. Its object is to render that treatise thoroughly intelligible. The author has aimed to illustrate, as preparatory to the detailed explanation of the work, the general bearings and relations of the Art of Rhetoric in itself, as well as the special mode of treating it adopted by Aristotle in his peculiar system. The evidence upon obscure or doubtful questions connected with the subject is examined; and the relations which Rhetoric bears, in Aristotle's view, to the kindred art of Logic are fully considered. A connected Analysis of the work is given, sometimes in the form of paraphrase; and a few important matters are separately discussed in Appendices. There is added, as a general Appendix, by way of specimen of the antagonistic system of Isocrates and others, a complete analysis of the treatise called Ῥητορικὴ πρὸς Ἀλέξανδρον, *with a discussion of its authorship and of the probable results of its teaching.*

CATULLI VERONENSIS LIBER. Edited by R. ELLIS, Fellow of Trinity College, Oxford. 18mo. 3*s.* 6*d.*

An elaborate Preface deals with the arrangement of the verse, and discusses various readings. "Rarely," says the Saturday Review, *"have we read a classic author with so reliable, acute, and safe a guide."*

Cicero.—THE SECOND PHILIPPIC ORATION. With an Introduction and Notes, translated from the German of KARL HALM. Edited, with Corrections and Additions, by JOHN E. B. MAYOR, M.A., Fellow and Classical Lecturer of St. John's College, Cambridge. Third Edition, revised. Fcap. 8vo. 5*s.*

This volume opens with a List of Books useful to the Student of Cicero, including History, Chronology, Lexicons, and some account of various editions, mostly German, of the works of Cicero. The Introduction is based on Halm: where Halm gives a reference to a classic, the passage has been commonly printed at length; where the reference is to Halm's notes on other Ciceronian speeches, or to modern books, the additional matter has been incorporated: and the numerous Greek quotations have been rendered into English. The English editor has further illustrated the work by additions drawn, for the most part, (1) *from the ancient authorities;* (2) *from his own private marginal references, and from collections;* (3) *from the notes of previous commentators. A copious 'argument' is also given.*

DEMOSTHENES ON THE CROWN. The Greek Text with English Notes. By B. DRAKE, M.A., late Fellow of King's College, Cambridge. Third Edition, to which is prefixed ÆSCHINES AGAINST CTESIPHON, with English Notes. Fcap. 8vo. 5*s.*

An Introduction discusses the immediate causes of the two orations, and their general character. The Notes contain frequent references to the best authorities. Among the appendices at the end is a chronological table of the life and public career of Æschines and Demosthenes.

Hodgson.—MYTHOLOGY FOR LATIN VERSIFICATION. A brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. HODGSON, B.D., late Provost of Eton. New Edition, revised by F. C. HODGSON, M.A. 18mo. 3*s.*

The late Provost of Eton has here supplied a help to the composition of Latin Verse, combined with a brief introduction to Classical Mythology.

In this new edition a few mistakes have been rectified; rules have been added to the Prosody; and a more uniform system has been adopted with regard to the help afforded.

JUVENAL FOR SCHOOLS. With English Notes. By J. E. B. MAYOR, M.A. New and cheaper Edition. Crown 8vo. [*In the Press.*

Marshall.—A TABLE OF IRREGULAR GREEK VERBS, classified according to the arrangement of Curtius' Greek Grammar. By J. M. MARSHALL, M.A., Fellow and late Lecturer of Brasenose College, Oxford; one of the Masters in Clifton College. 8vo. cloth. 1*s.*

The system of this table has been borrowed from the excellent Greek Grammar of Dr. Curtius.

Mayor, John E. B.—FIRST GREEK READER. Edited after KARL HALM, with Corrections and large Additions by JOHN E. B. MAYOR, M.A. Fellow and Classical Lecturer of St. John's College, Cambridge. Fcap. 8vo. 6*s.*

A selection of short passages, serving to illustrate especially the Greek Accidence. A good deal of syntax is incidentally taught, and Madvig and other books are cited, for the use of masters: but no learner *is expected to know more of syntax than is contained in the Notes and Vocabulary. A preface "To the Reader," not only explains the aim and method of the volume, but also deals with classical instruction generally. The extracts are uniformly in the Attic dialect, and any Hellenistic forms occurring in the original classic authors, such as Ælian and Polybius, have been discarded in favour of the corresponding Attic expressions. This book may be used in connexion with Mayor's "Greek for Beginners."*

Mayor, Joseph B.—GREEK FOR BEGINNERS. By the Rev. J. B. MAYOR, M.A. Fcap. 8vo. 4*s.* 6*d.*

The distinctive method of this book consists in building up a boy's knowledge of Greek upon the foundation of his knowledge of English and Latin, instead of trusting everything to the unassisted memory. The forms and constructions of Greek have been thoroughly compared with *those of Latin, and no Greek words have been used in the earlier part of the book except such as have connexions either in English or Latin. Each step leads naturally on to its successor, grammatical forms and rules are at once applied in a series of graduated exercises, accompanied by ample vocabularies. Thus the book serves as Grammar, Exercise book, and Vocabulary. Where possible, the Grammar has been simplified; the ordinary ten declensions are reduced to three, which correspond to the first three in Latin; and the system of* stems *is adopted. A general Vocabulary, and Index of Greek words, completes the work.*

Plato.—THE REPUBLIC OF PLATO. Translated into English, with an Analysis and Notes, by J. LL. DAVIES, M.A., and D. J. VAUGHAN, M.A. Third Edition, with Vignette Portraits of Plato and Socrates, engraved by JEENS from an Antique Gem. 18mo. 4*s*. 6*d*.

An introductory notice supplies some account of the life of Plato, and the translation is preceded by an elaborate analysis. "The translators have," in the judgment of the Saturday Review, *"produced a book which any reader, whether acquainted with the original or not, can peruse with pleasure as well as profit."*

Plautus (Ramsay).—THE MOSTELLARIA OF PLAUTUS. With Notes Critical and Explanatory, Prolegomena, and Excursus. By WILLIAM RAMSAY, M.A., formerly Professor of Humanity in the University of Glasgow. Edited by Professor GEORGE G. RAMSAY, M.A., of the University of Glasgow. 8vo. 14*s*.

"The fruits of that exhaustive research and that ripe and well-digested scholarship which its author brought to bear upon everything that he undertook are visible throughout it. It is furnished with a complete apparatus of prolegomena, notes, and excursus; and for the use of veteran scholars it probably leaves nothing to be desired."—PALL MALL GAZETTE.

Roby.—A LATIN GRAMMAR for the Higher Classes in Grammar Schools. By H. J. ROBY, M.A., based on the Elementary Latin Grammar. [*In the Press.*

Sallust.—CAII SALLUSTII CRISPI CATILINA ET JUGURTHA. For Use in Schools. With copious Notes. By C. MERIVALE, B.D. (In the present Edition the Notes have been carefully revised, and a few remarks and explanations added.) Second Edition. Fcap. 8vo. 4*s.* 6*d.*

The JUGURTHA and the CATILINA may be had separately, price 2*s.* 6*d.* each.

This edition of Sallust, prepared by the distinguished historian of Rome, contains an introduction, concerning the Life and works of Sallust, lists of the Consuls, and elaborate notes.

Tacitus.—THE HISTORY OF TACITUS TRANSLATED INTO ENGLISH. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Notes and a Map. 8vo. 10*s.* 6*d.*

The translators have endeavoured to adhere as closely to the original as was thought consistent with a proper observance of English idiom. At the same time, it has been their aim to reproduce the precise expressions of the author. The campaign of Civilis is elucidated in a note of some length, which is illustrated by a map, containing the names of places and of tribes occurring in the work. There is also a complete account of the Roman army as it was constituted in the time of Tacitus. This work is characterised by the Spectator *as "a scholarly and faithful translation."*

THE AGRICOLA AND GERMANIA OF TACITUS. A Revised Text, English Notes, and Maps. By ALFRED J. CHURCH, M.A., and W. J. BRODRIBB, M.A. Fcap. 8vo. 3*s.* 6*d.*

"We have endeavoured, with the aid of recent editions, thoroughly to elucidate the text, explaining the various difficulties, critical and grammatical, which occur to the student. We have consulted throughout, besides the older commentators, the editions of Ritter and Orelli, but we are under special obligations to the labours of the recent German editors, Wex, and Kritz." Two Indexes are appended, (1) *of Proper Names,* (2) *of Words and Phrases explained.*

THE AGRICOLA and GERMANIA may be had separately, price 2*s.* each.

Tacitus—*continued.*

THE AGRICOLA AND GERMANIA. Translated into English by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Maps and Notes. Extra fcap. 8vo. 2*s.* 6*d.*

The translators have sought to produce such a version as may satisfy scholars who demand a faithful rendering of the original, and English readers who are offended by the baldness and frigidity which commonly disfigure translations. The treatises are accompanied by introductions, notes, maps, and a chronological summary. The Athenæum *says of this work that it is "a version at once readable and exact, which may be perused with pleasure by all, and consulted with advantage by the classical student."*

Thring.—A LATIN GRADUAL. A First Latin Construing Book for Beginners. By EDWARD THRING, M.A. Fcap. 8vo. 2*s.* 6*d.*

The Head Master of Uppingham has here sought to supply by easy steps a knowledge of grammar, combined with a good Vocabulary. Passages have been selected from the best Latin authors in prose and verse. These passages are gradually built up in their grammatical structure, and finally printed in full. A short practical manual of common mood constructions, with their English equivalents, forms a second part.

A MANUAL OF MOOD CONSTRUCTIONS. Fcap. 8vo. 1*s.* 6*d.*

Treats of the ordinary mood constructions, as found in the Latin, Greek, and English languages.

A CONSTRUING BOOK. Fcap. 8vo. 2*s.* 6*d.*

Thucydides.—THE SICILIAN EXPEDITION. Being Books VI. and VII. of Thucydides, with Notes. A New Edition, revised and enlarged, with a Map. By the Rev. PERCIVAL FROST, M.A., late Fellow of St. John's College, Cambridge. Fcap. 8vo. 5*s.*

This edition is mainly a grammatical one. Attention is called to the force of compound verbs, and the exact meaning of the various tenses employed.

Wright.—Works by T. WRIGHT, M.A., late Head Master of Sutton Coldfield School.

HELLENICA; OR, A HISTORY OF GREECE IN GREEK, as related by Diodorus and Thucydides; being a First Greek Reading Book, with explanatory Notes, Critical and Historical. Third Edition, with a Vocabulary. 12mo. 3*s.* 6*d.*

In the last twenty chapters of this volume, Thucydides sketches the rise and progress of the Athenian Empire in so clear a style and in such simple language, that the editor has doubts whether any easier or more instructive passages can be selected for the use of the pupil who is commencing Greek. This book includes a chronological table of the events recorded.

A HELP TO LATIN GRAMMAR; or, The Form and Use of Words in Latin, with Progressive Exercises. Crown 8vo. 4*s.* 6*d.*

This book is not intended as a rival to any of the excellent Grammars now in use; but as a help to enable the beginner to understand them.

THE SEVEN KINGS OF ROME. An Easy Narrative, abridged from the First Book of Livy by the omission of Difficult Passages; being a First Latin Reading Book, with Grammatical Notes. With Vocabulary and Exercises. Second Edition. 5*s.*

This work is intended to supply the pupil with an easy construing book, which may at the same time be made the vehicle for instructing him in the rules of grammar and principles of composition. The notes profess to teach what is commonly taught in grammars. It is conceived that the pupil will learn the rules of construction of the language much more easily from separate examples, which are pointed out to him in the course of his reading, and which he may himself set down in his note-book after some scheme of his own, than from a heap of quotations amassed for him by others.

Or, separately,

SEVEN KINGS OF ROME. 3*s.*

VOCABULARY AND EXERCISES TO "THE SEVEN KINGS." 2*s.* 6*d.*

CLASSIC VERSIONS OF ENGLISH BOOKS, AND LATIN HYMNS.

THE following works are, as the heading indicates, classic renderings of English books. For scholars, and particularly for writers of Latin Verse, the series has a special value. The Hymni Ecclesiæ are here inserted, as partly falling under the same class.

Church and Kebbel.—HORÆ TENNYSONIANÆ. Curantibus A. J. CHURCH, A.M., T. E. KEBBEL, A.M. [*In the Press.*

Latham.—SERTUM SHAKSPERIANUM, Subnexis aliquot aliunde excerptis floribus. Latine reddidit Rev. H. LATHAM, M.A. Extra fcap. 8vo. 5*s.*

Besides versions of Shakspeare this volume contains, among other pieces, Gray's "Elegy," Campbell's "Hohenlinden," Wolfe's "Burial of Sir John Moore," and selections from Cowper and George Herbert.

Lyttelton.—THE COMUS OF MILTON, rendered into Greek Verse. By LORD LYTTELTON. Extra fcap. 8vo. 6*s.* 6*d.*

THE SAMSON AGONISTES OF MILTON, rendered into Greek Verse. By LORD LYTTELTON. Extra fcap. 8vo. 6*s.* 6*d.*

Merivale.—KEATS' HYPERION, rendered into Latin Verse. By C. MERIVALE, B.D. Second Edit. Extra fcap. 8vo. 3*s.* 6*d.*

Hymni Ecclesiæ.—Edited by Rev. DR. NEWMAN. Extra fcap. 8vo. 7*s.* 6*d.*

Hymns of the Mediæval Church. The first Part contains selections from the Parisian Breviary; the second from those of Rome, Salisbury, and York.

Trench (Archbishop).—SACRED LATIN POETRY, chiefly Lyrical, selected and arranged for Use; with Notes and Introduction. Fcap. 8vo. 7*s.*

In this work the editor has selected hymns of a catholic religious sentiment that are common to Christendom, while rejecting those of a distinctively Romish character.

MATHEMATICS.

Airy.—Works by G. B. AIRY, Astronomer Royal :—

ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS. Designed for the Use of Students in the Universities. With Diagrams. Crown 8vo. cloth. 5*s*. 6*d*.

It is hoped that the methods of solution here explained, and the instances exhibited, will be found sufficient for application to nearly all the important problems of Physical Science, which require for their complete investigation the aid of Partial Differential Equations.

ON THE ALGEBRAICAL AND NUMERICAL THEORY OF ERRORS OF OBSERVATIONS AND THE COMBINATION OF OBSERVATIONS. Crown 8vo. cloth. 6*s*. 6*d*.

In order to spare astronomers and observers in natural philosophy the confusion and loss of time which are produced by referring to the ordinary treatises embracing both *branches of probabilities (the first relating to chances which can be altered only by the changes of entire units or integral multiples of units in the fundamental conditions of the problem; the other concerning those chances which have respect to insensible gradations in the value of the element measured) the present tract has been drawn up. It relates only to errors of observation, and to the rules, derivable from the consideration of these errors, for the combination of the results of observations.*

Airy, G. B.—*continued.*

UNDULATORY THEORY OF OPTICS. Designed for the Use of Students in the University. New Edition. Crown 8vo. cloth. 6*s.* 6*d.*

The undulatory theory of optics is presented to the reader as having the same claims to his attention as the theory of gravitation: namely, that it is certainly true, and that, by mathematical operations of general elegance, it leads to results of great interest. This theory explains with accuracy a vast variety of phenomena of the most complicated kind. The plan of this tract has been to include those phenomena only which admit of calculation, and the investigations are applied only to phenomena which actually have been observed.

ON SOUND AND ATMOSPHERIC VIBRATIONS. With the Mathematical Elements of Music. Designed for the Use of Students of the University. Crown 8vo. 9*s.*

This volume consists of sections, which again are divided into numbered articles, on the following topics: General recognition of the air as the medium which conveys sound; Properties of the air on which the formation and transmission of sound depend; Theory of undulations as applied to sound, &c.; Investigation of the motion of a wave of air through the atmosphere; Transmission of waves of soniferous vibrations through different gases, solids, and fluids; Experiments on the velocity of sound, &c.; On musical sounds, and the manner of producing them; On the elements of musical harmony and melody, and of simple musical composition; On instrumental music; On the human organs of speech and hearing.

Bayma.—THE ELEMENTS OF MOLECULAR MECHANICS. By JOSEPH BAYMA, S. J., Professor of Philosophy, Stonyhurst College. Demy 8vo. cloth. 10*s.* 6*d.*

Of the twelve Books into which the present treatise is divided, the first and second give the demonstration of the principles which bear directly on the constitution and the properties of matter. The next three books contain a series of theorems and of problems on the laws of motion of elementary substances. In the sixth and seventh, the mechanical constitution of mole-

cules is investigated and determined: and by it the general properties of bodies are explained. The eighth book treats of luminiferous æther. The ninth explains some special properties of bodies. The tenth and eleventh contain a radical and lengthy investigation of chemical principles and relations, which may lead to practical results of high importance. The twelfth and last book treats of molecular masses, distances, and powers.

Beasley.—AN ELEMENTARY TREATISE ON PLANE TRIGONOMETRY. With Examples. By R. D. BEASLEY, M.A., Head Master of Grantham Grammar School. Second Edition, revised and enlarged. Crown 8vo. cloth. 3*s.* 6*d.*

This treatise is specially intended for use in schools. The choice of matter has been chiefly guided by the requirements of the three days' examination at Cambridge. About four hundred examples have been added to this edition, mainly collected from the examination papers of the last ten years.

Boole.—Works by G. BOOLE, D.C.L., F.R.S., Professor of Mathematics in the Queen's University, Ireland.

A TREATISE ON DIFFERENTIAL EQUATIONS. New and Revised Edition. Edited by I. TODHUNTER. Crown 8vo. cloth. 14*s.*

Professor Boole has endeavoured in this treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations, as was consistent with the idea of a work intended, primarily, for elementary instruction. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable for the beginner, while the latter ones are devoted either to an account of recent discovery, or the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connexion with the methods and processes of his previous course.

A TREATISE ON DIFFERENTIAL EQUATIONS. Supplementary Volume. Edited by I. TODHUNTER. Crown 8vo. cloth. 8*s.* 6*d.*

This volume contains all that Professor Boole wrote for the purpose of enlarging his treatise on Differential Equations.

Boole, G., D.C.L.—*continued.*

THE CALCULUS OF FINITE DIFFERENCES. Crown 8vo. cloth. 10*s.* 6*d.*

In this exposition of the Calculus of Finite Differences, particular attention has been paid to the connexion of its methods with those of the Differential Calculus—a connexion which in some instances involves far more than a merely formal analogy. The work is in some measure designed as a sequel to Professor Boole's Treatise on Differential Equations.

CAMBRIDGE SENATE-HOUSE PROBLEMS AND RIDERS, WITH SOLUTIONS:—

1848-1851.—PROBLEMS. By FERRERS and JACKSON. 8vo. cloth. 15*s.* 6*d.*

1848-1851.—RIDERS. By JAMESON. 8vo. cloth. 7*s.* 6*d.*

1854.—PROBLEMS AND RIDERS. By WALTON and MACKENZIE. 8vo. cloth. 10*s.* 6*d.*

1857.—PROBLEMS AND RIDERS. By CAMPION and WALTON. 8vo. cloth. 8*s.* 6*d.*

1860.—PROBLEMS AND RIDERS. By WATSON and ROUTH. Crown 8vo. cloth. 7*s.* 6*d.*

1864.—PROBLEMS AND RIDERS. By WALTON and WILKINSON. 8vo. cloth. 10*s.* 6*d.*

These volumes will be found of great value to Teachers and Students, as indicating the style and range of mathematical study in the University of Cambridge.

CAMBRIDGE COURSE OF ELEMENTARY NATURAL PHILOSOPHY, for the Degree of B.A. Originally compiled by J. C. SNOWBALL, M.A., late Fellow of St. John's College. Fifth Edition, revised and enlarged, and adapted for the Middle-Class Examinations by THOMAS LUND, B.D., Late Fellow and Lecturer of St. John's College, Editor of Wood's Algebra, &c. Crown 8vo. cloth. 5*s.*

This work will be found adapted to the wants, not only of University Students, but also of many others who require a short course of Mechanics *and Hydrostatics, and especially of the candidates at our Middle Class Examinations. At the end of each chapter a series of easy questions is added for the exercise of the student.*

CAMBRIDGE AND DUBLIN MATHEMATICAL JOURNAL. The Complete Work, in Nine Vols. 8vo. cloth, 7*l.* 4*s.*

Only a few copies remain on hand. Among Contributors to this work will be found Sir W. Thomson, Stokes, Adams, Boole, Sir W. R. Hamilton, De Morgan, Cayley, Sylvester, Jellett, and other distinguished mathematicians.

Candler.—HELP TO ARITHMETIC. Designed for the use of Schools. By H. CANDLER, M.A. Mathematical Master of Uppingham School. Extra fcap. 8vo. 2*s.* 6*d.*

This work is intended as a companion to any text book that may be in use.

Cheyne.—AN ELEMENTARY TREATISE ON THE PLANETARY THEORY. With a Collection of Problems. By C. H. H. CHEYNE, B.A. Crown 8vo. cloth. 6*s.* 6*d.*

In this volume, an attempt has been made to produce a treatise on the Planetary theory, which, being elementary in character, should be so far complete, as to contain all that is usually required by students in the University of Cambridge.

THE EARTH'S MOTION OF ROTATION. By C. H. H. CHEYNE, M.A. Crown 8vo. 3*s.* 6*d.*

The first part of this work consists of an application of the method of the variation of elements to the general problem of rotation. In the second part the general rotation formulæ are applied to the particular case of the earth.

Childe.—THE SINGULAR PROPERTIES OF THE ELLIPSOID AND ASSOCIATED SURFACES OF THE NTH DEGREE. By the Rev. G. F. CHILDE, M.A., Author of "Ray Surfaces," "Related Caustics," &c. 8vo. 10*s*. 6*d*.

The object of this volume is to develop peculiarities in the Ellipsoid; and, further, to establish analagous properties in the unlimited congeneric series of which this remarkable surface is a constituent.

Christie.—A COLLECTION OF ELEMENTARY TEST-QUESTIONS IN PURE AND MIXED MATHEMATICS; with Answers and Appendices on Synthetic Division, and on the Solution of Numerical Equations by Horner's Method. By JAMES R. CHRISTIE, F.R.S., late First Mathematical Master at the Royal Military Academy, Woolwich. Crown 8vo. cloth. 8*s*. 6*d*.

The series of Mathematical exercises here offered to the public is collected from those which the author has, from time to time, proposed for solution by his pupils during a long career at the Royal Military Academy. A student who finds that he is able to solve the larger portion of these exercises, may consider that he is thoroughly well grounded in the elementary principles of pure and mixed Mathematics.

Dalton.—ARITHMETICAL EXAMPLES. Progressively arranged, with Exercises and Examination Papers. By the Rev. T. DALTON, M.A., Assistant Master of Eton College. 18mo. cloth. 2*s*. 6*d*.

Answers to the Examples are appended.

Day.—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY. PART I., THE ELLIPSE, with Problems. By the Rev. H. G. DAY, M.A., Head Master of Sedburgh Grammar School. Crown 8vo. 3*s*. 6*d*.

The object of this book is the introduction of a treatment of Conic Sections which should be simple and natural, and lead by an easy transition to the analytical methods, without departing from the strict geometry of Euclid.

Dodgson.—AN ELEMENTARY TREATISE ON DETERMINANTS, with their Application to Simultaneous Linear Equations and Algebraical Geometry. By CHARLES L. DODGSON, M.A., Student and Mathematical Lecturer of Christ Church, Oxford. Small 4to. cloth. 10*s.* 6*d.*

The object of the author is to present the subject as a continuous chain of argument, separated from all accessories of explanation or illustration. All such explanation and illustration as seemed necessary for a beginner, are introduced either in the form of foot-notes, or, where that would have occupied too much room, of Appendices.

Drew.—GEOMETRICAL TREATISE ON CONIC SECTIONS. By W. H. DREW, M.A., St. John's College, Cambridge. Third Edition. Crown 8vo. cloth. 4*s.* 6*d.*

In this work the subject of Conic Sections has been placed before the student in such a form that, it is hoped, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies. With a view, also, of rendering the work a complete manual of what is required at the Universities, there have either been embodied into the text or inserted among the examples, every book-work question, problem, and rider, which has been proposed in the Cambridge examinations up to the present time.

SOLUTIONS TO THE PROBLEMS IN DREW'S CONIC SECTIONS. Crown 8vo. cloth. 4*s.* 6*d.*

Ferrers.—AN ELEMENTARY TREATISE ON TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. By the Rev. N. M. FERRERS, M.A., Fellow and Tutor of Gonville and Caius College, Cambridge. Second Edition. Crown 8vo. 6*s.* 6*d.*

The object of the author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian system, instead of regarding it as only a special form of Abridged Notation. A short chapter on Determinants has been introduced.

Frost.—THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a collection of Problems, principally intended as Examples of Newton's Methods. By PERCIVAL FROST, M.A., late Fellow of St. John's College, Mathematical Lecturer of King's College, Cambridge. Second Edition. 8vo. cloth. 10*s*. 6*d*.

The author's principal intention is to explain difficulties which may be encountered by the student on first reading the Principia, and to illustrate the advantages of a careful study of the methods employed by Newton, by showing the extent to which they may be applied in the solution of problems; he has also endeavoured to give assistance to the student who is engaged in the study of the higher branches of mathematics, by representing in a geometrical form several of the processes employed in the Differential and Integral Calculus, and in the analytical investigations of Dynamics.

Frost and Wolstenholme.—A TREATISE ON SOLID GEOMETRY. By PERCIVAL FROST, M.A., and the Rev. J. WOLSTENHOLME, M.A., Fellow and Assistant Tutor of Christ's College. 8vo. cloth. 18*s*.

The authors have endeavoured to present before students as comprehensive a view of the subject as possible. Intending to make the subject accessible, at least in the earlier portion, to all classes of students, they have endeavoured to explain completely all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have considered themselves to be addressing a higher class of students; and they have there tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.

Godfray.—A TREATISE ON ASTRONOMY, for the Use of Colleges and Schools. By HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge. 8vo. cloth. 12*s*. 6*d*.

This book embraces all those branches of Astronomy which have, from time to time, been recommended by the Cambridge Board of Mathematical

Godfray, Hugh—*continued.*

Studies: but by far the larger and easier portion, adapted to the first three days of the Examination for Honours, may be read by the more advanced pupils in many of our schools. The author's aim has been to convey clear and distinct ideas of the celestial phenomena.

AN ELEMENTARY TREATISE ON THE LUNAR THEORY, with a Brief Sketch of the Problem up to the time of Newton. By HUGH GODFRAY, M.A. Second Edition, revised. Crown 8vo. cloth. 5*s.* 6*d.*

These pages will, it is hoped, form an introduction to more recondite works. Difficulties have been discussed at considerable length. The selection of the method followed with regard to analytical solutions, which is the same as that of Airy, Herschel, &c. was made on account of its simplicity; it is, moreover, the method which has obtained in the University of Cambridge.

Hemming.—AN ELEMENTARY TREATISE ON THE DIFFERENTIAL AND INTEGRAL CALCULUS, for the Use of Colleges and Schools. By G. W. HEMMING, M.A., Fellow of St. John's College, Cambridge. Second Edition, with Corrections and Additions. 8vo. cloth. 9*s.*

Jones and Cheyne.—ALGEBRAICAL EXERCISES. Progressively arranged. By the Rev. C. A. JONES, M.A., and C. H. CHEYNE, M.A., Mathematical Masters of Westminster School. New Edition. 18mo. cloth. 2*s.* 6*d.*

This little book is intended to meet a difficulty which is probably felt more or less by all engaged in teaching Algebra to beginners. It is, that while new ideas are being acquired, old ones are forgotten. In the belief that constant practice is the only remedy for this, the present series of miscellaneous exercises has been prepared. Their peculiarity consists in this, that though miscellaneous they are yet progressive, and may be used by *the pupil almost from the commencement of his studies. They are not intended to supersede the systematically arranged examples to be found in ordinary treatises on Algebra, but rather to supplement them.*

The book being intended chiefly for Schools and Junior Students, the higher parts of Algebra have not been included.

Kitchener.—A GEOMETRICAL NOTE-BOOK, containing Easy Problems in Geometrical Drawing preparatory to the Study of Geometry. For the Use of Schools. By F. E. KITCHENER, M.A., Mathematical Master at Rugby. 4to. 2*s.*

It is the object of this book to make some way in overcoming the difficulties of Geometrical conception, before the mind is called to the attack of Geometrical theorems. A few simple methods of construction are given; and space is left on each page, in order that the learner may draw in the figures.

Morgan.—A COLLECTION OF PROBLEMS AND EXAMPLES IN MATHEMATICS. With Answers. By H. A. MORGAN, M.A., Sadlerian and Mathematical Lecturer of Jesus College, Cambridge. Crown 8vo. cloth. 6*s.* 6*d.*

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the papers set during late years at Jesus College. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

Parkinson.—Works by S. PARKINSON, B.D., Fellow and Prælector of St. John's College, Cambridge.

AN ELEMENTARY TREATISE ON MECHANICS. For the Use of the Junior Classes at the University and the Higher Classes in Schools. With a Collection of Examples. Fourth edition, revised. Crown 8vo. cloth. 9*s.* 6*d.*

In preparing a fourth edition of this work the author has kept the same object in view as he had in the former editions—namely, to include in it such portions of Theoretical Mechanics as can be conveniently investigated

Parkinson, S.—*continued.*

without the use of the Differential Calculus, and so render it suitable as a manual for the junior classes in the University and the higher classes in Schools. With one or two short exceptions, the student is not presumed to require a knowledge of any branches of Mathematics beyond the elements of Algebra, Geometry, and Trigonometry. Several additional propositions have been incorporated in the work for the purpose of rendering it more complete; and the collection of Examples and Problems has been largely increased.

A TREATISE ON OPTICS. Second Edition, revised. Crown 8vo. cloth. 10*s.* 6*d.*

A collection of examples and problems has been appended to this work, which are sufficiently numerous and varied in character to afford useful exercise for the student. For the greater part of them, recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

Phear.—ELEMENTARY HYDROSTATICS. With Numerous Examples. By J. B. PHEAR, M.A., Fellow and late Assistant Tutor of Clare College, Cambridge. Fourth Edition. Crown 8vo. cloth. 5*s.* 6*d.*

This edition has been carefully revised throughout, and many new illustrations and examples added, which it is hoped will increase its usefulness to students at the Universities and in Schools. In accordance with suggestions from many engaged in tuition, answers to all the Examples have been given at the end of the book.

Pratt.—A TREATISE ON ATTRACTIONS, LAPLACE'S FUNCTIONS, AND THE FIGURE OF THE EARTH. By JOHN H. PRATT, M.A., Archdeacon of Calcutta, Author of "The Mathematical Principles of Mechanical Philosophy." Third Edition. Crown 8vo. cloth. 6*s.* 6*d.*

The author's chief design in this treatise is to give an answer to the question, "Has the Earth acquired its present form from being originally in a fluid state?" This Edition is a complete revision of the former ones.

Puckle.—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. With Numerous Examples and Hints for their Solution; especially designed for the Use of Beginners. By G. H. PUCKLE, M.A., Head Master of Windermere College. Third Edition, revised and enlarged. Crown 8vo. cloth. 7*s*. 6*d*.

This work has been completely rewritten, and a large amount of new matter has been added to suit the requirements of the present time. The author has written with special reference to those difficulties and misapprehensions which are found most common to beginners. The treatise is complete as far as regards Cartesian Co-ordinates.

Rawlinson.—ELEMENTARY STATICS, by the Rev. GEORGE RAWLINSON, M.A. Edited by the Rev. EDWARD STURGES, M.A., of Emmanuel College, Cambridge, and late Professor of the Applied Sciences, Elphinstone College, Bombay. Crown 8vo. cloth. 4*s*. 6*d*.

Published under the authority of Her Majesty's Secretary of State for India, for use in the Government Schools and Colleges in India.

Reynolds.—MODERN METHODS IN ELEMENTARY GEOMETRY. By E. M. REYNOLDS, M.A., Mathematical Master in Clifton College. Crown 8vo. 3*s*. 6*d*.

Geometry has received extensive developments in modern times, but in England there has been no corresponding improvement in elementary teaching.

Routh.—AN ELEMENTARY TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With Numerous Examples. By EDWARD JOHN ROUTH, M.A., late Fellow and Assistant Tutor of St. Peter's College, Cambridge; Examiner in the University of London. Second Edition, enlarged. Crown 8vo. cloth. 14*s*.

In this edition the author has made several additions to each chapter. He has tried, even at the risk of some little repetition, to make each *chapter, as far as possible, complete in itself, so that all that relates to any one part of the subject may be found in the same place. This arrangement will enable every student to select his own order in which to read the subject. The Examples which will be found at the end of each chapter have been chiefly selected from the Examination Papers which have been set in the University and the Colleges in the last few years.*

Smith (Barnard).—Works by BARNARD SMITH, M.A., Rector of Glaston, Rutlandshire, late Fellow and Senior Bursar of St. Peter's College, Cambridge.

ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. Tenth Edition. Crown 8vo. cloth. 10*s.* 6*d.*

This manual is now extensively used in Schools and Colleges, both in England and in the Colonies. It has also been found of great service for students preparing for the Middle Class and Civil and Military Service Examinations, from the care that has been taken to elucidate the principles *of all the rules. The present edition has been carefully revised. "To all those whose minds are sufficiently developed to comprehend the simplest mathematical reasoning, and who have not yet thoroughly mastered the principles of Arithmetic and Algebra, it is calculated to be of great advantage."*—ATHENÆUM.

Of this work, also, one of the highest possible authorities, the late Dean Peacock, writes: "Mr. Smith's work is a most useful publication. The rules are stated with great clearness. The examples are well selected, and worked out with just sufficient detail, without being encumbered by too minute explanations; and there prevails throughout it that just proportion of theory and practice, which is the crowning excellence of an elementary work."

Smith, Barnard—*continued.*

ARITHMETIC FOR SCHOOLS. New Edition. Crown 8vo. cloth. 4*s.* 6*d.*

Adapted from the author's work on "Arithmetic and Algebra," by the omission of the algebraic portion, and by the introduction of new exercises. The reason of each arithmetical process is fully exhibited. The system of Decimal Coinage is explained; and answers to the exercises are appended at the end. This Arithmetic is characterised as "admirably adapted for instruction, combining just sufficient theory with a large and well-selected collection of exercises for practice."—JOURNAL OF EDUCATION.

COMPANION TO ARITHMETIC FOR SCHOOLS. [*Preparing.*

A KEY TO THE ARITHMETIC FOR SCHOOLS. Seventh Edition. Crown 8vo. cloth. 8*s.* 6*d.*

EXERCISES IN ARITHMETIC. With Answers. Crown 8vo. limp cloth. 2*s.* 6*d.*

Or sold separately, Part I. 1*s.*; Part II. 1*s.*; Answers, 6*d.*

These Exercises have been published in order to give the pupil examples in every rule of Arithmetic. The greater number have been carefully compiled from the latest University and School Examination Papers.

SCHOOL CLASS-BOOK OF ARITHMETIC. 18mo. cloth. 3*s.*

Or sold separately, Parts I. and II. 10*d.* each; Part III. 1*s.*

This manual, published at the request of many schoolmasters, and chiefly intended for National and Elementary Schools, has been prepared on the same plan as that adopted in the author's School Arithmetic, which is in extensive circulation in England and abroad. The Metrical Tables have been introduced, from the conviction on the part of the author, that the knowledge of such tables, and the mode of applying them, will be of great use to the rising generation.

KEYS TO SCHOOL CLASS-BOOK OF ARITHMETIC. Complete in one volume, 18mo. cloth, 6*s.* 6*d.*; or Parts I. II. and III. 2*s.* 6*d.* each.

Smith, Barnard—*continued.*

SHILLING BOOK OF ARITHMETIC FOR NATIONAL AND ELEMENTARY SCHOOLS. 18mo. cloth. Or separately, Part I. 2*d.*; Part II. 3*d.*; Part III. 7*d.* Answers, 6*d.*

THE SAME, with Answers complete. 18mo. cloth. 1*s.* 6*d.*

This Shilling Book of Arithmetic has been prepared for the use of National and other schools at the urgent request of numerous masters of schools both at home and abroad. The Explanations of the Rules, and the Examples will, it is hoped, be found suited to the most elementary classes.

KEY TO SHILLING BOOK OF ARITHMETIC. 18mo. cloth. 4*s.* 6*d.*

EXAMINATION PAPERS IN ARITHMETIC. 18mo. cloth. 1*s.* 6*d.* The same, with Answers, 18mo. 1*s.* 9*d.*

The object of these Examination Papers is to test students both in the theory and practice of Arithmetic. It is hoped that the method adopted will lead students to deduce results from general principles rather than to apply stated rules. The author believes that the practice of giving examples under particular rules makes the working of Arithmetic quite mechanical, and tends to throw all but very clever boys off their balance when a general paper on the subject is put before them.

KEY TO EXAMINATION PAPERS IN ARITHMETIC. 18mo. cloth. 4*s.* 6*d.*

Smith (J. H.).—Works by J. H. SMITH, M.A., Gonville and Caius College, Cambridge.

A TREATISE ON ELEMENTARY STATICS. Second Edition. 8vo. 5*s.* 6*d.*

This treatise is intended to give a simple explanation of the part of Statics required in the previous examination and the second examination for ordinary degrees in the University of Cambridge. The examples have been

Smith, J. H.—*continued.*

selected from papers set in University Examinations. In this edition some explanations of the elementary definitions have been added; and a new and simple proof of the first part of the "Parallelogram of Forces," a new introduction to the Principle of Moments, and a short account of Friction have been inserted.

A TREATISE ON ELEMENTARY HYDROSTATICS. 8vo. 4*s*. 6*d.*

An explanation of the subject of Elementary Hydrostatics as required in the general examination for ordinary degrees at Cambridge. The Examples given at the end of the book are taken from Papers set in the University Examinations.

A TREATISE ON ELEMENTARY TRIGONOMETRY. 8vo. 5*s*.

This work treats of the elementary parts of Trigonometry, as required of the Candidates for Honours in the previous examination.

A TREATISE ON ELEMENTARY ALGEBRA. For the Use ot Colleges and Schools. Crown 8vo. cloth. 6*s*. 6*d.*

The object of this treatise is to explain those parts of Algebra which precede the Binomial Theorem. The Author has carefully abstained from making extracts from works on the subject in common use. He has written at considerable length on the subject of Factors and Fractions, because those branches of the subject present great difficulties to beginners. The rule has been never to evade a difficulty. The Examples are progressive, easy, and to a great extent original. Some have been selected from the Senate House Papers of the last three years, and a few from old English, French, or German works.

Snowball.—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY; with the Construction and Use of Tables of Logarithms. By J. C. SNOWBALL, M.A. Tenth Edition. Crown 8vo. cloth. 7*s*. 6*d.*

In preparing the present edition for the press, the text has been subjected to a careful revision; the proofs of some of the more impor-

tant propositions have been rendered more strict and general; and a considerable addition of more than two hundred examples, taken principally from the questions set of late years in the public examinations of the University and of individual Colleges, has been made to the collection of Examples and Problems for practice.

Tait and Steele.—DYNAMICS OF A PARTICLE. With numerous Examples. By Professor TAIT and Mr. STEELE. New Edition. Crown 8vo. cloth. 10*s*. 6*d*.

In this treatise will be found all the ordinary propositions, connected with the Dynamics of Particles, which can be conveniently deduced without the use of D'Alembert's Principle. Throughout the book will be found a number of illustrative examples introduced in the text, and for the most part completely worked out; others with occasional solutions or hints to assist the student are appended to each chapter. For by far the greater portion of these, the Cambridge Senate-House and College Examination Papers have been applied to.

Taylor.—GEOMETRICAL CONICS; including Anharmonic Ratio and Projection, with numerous Examples. By C. TAYLOR, B.A., Scholar of St. John's College, Cambridge. Crown 8vo. cloth. 7*s*. 6*d*.

This work contains elementary proofs of the principal properties of Conic Sections, together with chapters on Projection and Anharmonic Ratio.

Tebay.—ELEMENTARY MENSURATION FOR SCHOOLS. With numerous Examples. By SEPTIMUS TEBAY, B.A., Head Master of Queen Elizabeth's Grammar School, Rivington. Extra fcap. 8vo. 3*s*. 6*d*.

The object of the present work is to enable boys to acquire a moderate knowledge of Mensuration in a reasonable time. All difficult and useless matter has been avoided. The examples for the most part are easy, and the rules are concise.

Todhunter.—Works by I. TODHUNTER, M.A., F.R.S., of St. John's College, Cambridge.

THE ELEMENTS OF EUCLID. For the Use of Colleges and Schools. New Edition. 18mo. cloth. 3*s.* 6*d.*

As the elements of Euclid are usually placed in the hands of young students, it is important to exhibit the work in such a form as will assist them in overcoming the difficulties which they experience on their first introduction to processes of continuous argument. No method appears to be so useful as that of breaking up the demonstrations into their constituent parts; a plan strongly recommended by Professor De Morgan. In the present Edition each distinct assertion in the argument begins a new line: and at the ends of the lines are placed the necessary references to the preceding principles on which the assertions depend. The longer propositions are distributed into subordinate parts, which are distinguished by breaks at the beginning of the lines. Notes, appendix, and a collection of exercises are added.

MENSURATION FOR BEGINNERS. With Numerous Examples. 18mo. cloth. 2*s.* 6*d.*

The subjects included in the present work are those which have usually found a place in Elementary Treatises on Mensuration. The mode of treatment has been determined by the fact that the work is intended for the use of beginners. Accordingly it is divided into short independent chapters, which are followed by appropriate examples. A knowledge of the elements of Arithmetic is all that is assumed; and in connection with most of the Rules of Mensuration it has been found practicable to give such explanations and illustrations as will supply the place of formal mathematical demonstrations, which would have been unsuitable to the character of the work.

ALGEBRA FOR BEGINNERS. With numerous Examples. New Edition. 18mo. cloth. 2*s.* 6*d.*

Great pains have been taken to render this work intelligible to young students, by the use of simple language and by copious explanations. In

Todhunter, I.—*continued.*

determining the subjects to be included and the space to be assigned to each, the Author has been guided by the papers given at the various examinations in elementary Algebra which are now carried on in this country. The book may be said to consist of three parts. The first part contains the elementary operations in integral and fractional expressions; the second the solution of equations and problems; the third treats of various subjects which are introduced but rarely into examination papers, and are more briefly discussed. Provision has at the same time been made for the introduction of easy equations and problems at an early stage—for those who prefer such a course.

KEY TO ALGEBRA FOR BEGINNERS. Crown 8vo. cloth. 6*s*. 6*d*.

TRIGONOMETRY FOR BEGINNERS. With numerous Examples. New Edition. 18mo. cloth. 2*s*. 6*d*.

Intended to serve as an introduction to the larger treatise on Plane Trigonometry, published by the Author. The same plan has been adopted as in the Algebra for Beginners: *the subject is discussed in short chapters, and a collection of examples is attached to each chapter. The first fourteen chapters present the geometrical part of Plane Trigonometry; and contain all that is necessary for practical purposes. The range of matter included is such as seems required by the various examinations in elementary Trigonometry which are now carried on in the country. Answers are appended at the end.*

MECHANICS FOR BEGINNERS. With numerous Examples. 18mo. cloth. 4*s*. 6*d*.

Intended as a companion to the two preceding books. The work forms an elementary treatise on demonstrative mechanics. It may be true that this part of mixed mathematics has been sometimes made too abstract and speculative; but it can hardly be doubted that a knowledge of the elements at least of the theory of the subject is extremely valuable even for those who are mainly concerned with practical results. The Author has accordingly endeavoured to provide a suitable introduction to the study of applied

Todhunter, I.—*continued.*

as well as of theoretical mechanics. The work consists of two parts, namely, Statics and Dynamics. It will be found to contain all that is usually comprised in elementary treatises on Mechanics, together with some additions.

ALGEBRA. For the Use of Colleges and Schools. Fourth Edition. Crown 8vo. cloth. 7*s*. 6*d*.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of Examples for Exercise. *The author has sought to render the work easily intelligible to students, without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples, about* Sixteen hundred and fifty *in number, have been selected with a view to illustrate every part of the subject. Each chapter is complete in itself; and the work will be found peculiarly adapted to the wants of students who are without the aid of a teacher. The Answers to the examples, with hints for the solution of some in which assistance may be needed, are given at the end of the book.*

AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS. Second Edition, revised. Crown 8vo. cloth. 7*s*. 6*d*.

This treatise contains all the propositions which are usually included in elementary treatises on the theory of Equations, together with Examples for exercise. These have been selected from the College and University Examination Papers, and the results have been given when it appeared necessary. In order to exhibit a comprehensive view of the subject, the treatise includes investigations which are not found in all the preceding elementary treatises, and also some investigations which are not to be found in any of them. For the second edition the work has been revised and some additions have been made, the most important being an account of the researches of Professor Sylvester respecting Newton's Rule.

PLANE TRIGONOMETRY. For Schools and Colleges. Third Edition. Crown 8vo. cloth. 5*s*.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of

Todhunter, I.—*continued.*

obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples: those which are entitled Miscellaneous Examples, *together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. In the Second Edition the hints for the solution of the Examples have been considerably increased.*

A TREATISE ON SPHERICAL TRIGONOMETRY. Second Edition, enlarged. Crown 8vo. cloth. 4*s.* 6*d.*

The present work is constructed on the same plan as the treatise on Plane Trigonometry, to which it is intended as a sequel. In the account of Napier's Rules of Circular Parts, an explanation has been given of a method of proof devised by Napier, which seems to have been overlooked by most modern writers on the subject. Considerable labour has been bestowed on the text in order to render it comprehensive and accurate, and the Examples (selected chiefly from College Examination Papers) have all been carefully verified.

PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. With numerous Examples. Fourth Edition, revised and enlarged. Crown 8vo. cloth. 7*s.* 6*d.*

The Author has here endeavoured to exhibit the subject in a simple manner for the benefit of beginners, and at the same time to include in one volume all that students usually require. In addition, therefore, to the propositions which have always appeared in such treatises, he has introduced the methods of abridged notation, *which are of more recent origin; these methods, which are of a less elementary character than the rest of the work, are placed in separate chapters, and may be ommitted by the student at first.*

A TREATISE ON THE DIFFERENTIAL CALCULUS. With numerous Examples. Fourth Edition. Crown 8vo. cloth. 10*s.* 6*d.*

The Author has endeavoured in the present work to exhibit a comprehensive view of the Differential Calculus on the method of limits. In the more elementary portions he has entered into considerable detail in the

Todhunter, I.—*continued.*

explanations, with the hope that a reader who is without the assistance of a tutor may be enabled to acquire a competent acquaintance with the subject. The method adopted is that of Differential Coefficients. To the different chapters are appended examples sufficiently numerous to render another book unnecessary; these examples being mostly selected from College Examination Papers.

A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS. With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10*s*. 6*d*.

This is designed as a work at once elementary and complete, adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, it has been sought to exhibit the principles clearly, and to illustrate all their most important results. The process of summation has been repeatedly brought forward, with the view of securing the attention of the student to the notions which form the true foundation of the Calculus itself, as well as of its most valuable applications. Every attempt has been made to explain those difficulties which usually perplex beginners, especially with reference to the limits *of integrations. A new method has been adopted in regard to the transformation of multiple integrals. The last chapter deals with the Calculus of Variations. A large collection of exercises, selected from College Examination Papers, has been appended to the several chapters.*

EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. Second Edition, revised. Crown 8vo. cloth. 4*s*.

A TREATISE ON ANALYTICAL STATICS. With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10*s*. 6*d*.

In this work on statics (treating of the laws of the equilibrium of bodies) will be found all the propositions which usually appear in treatises on Theoretical Statics. To the different chapters examples are appended, which have been principally selected from University Examination Papers.

In the Third Edition many additions have been made, in order to illustrate the application of the principles of the subject to the solution of problems.

Wilson, J. M. — ELEMENTARY GEOMETRY. Angles, Parallels, Triangles, and Equivalent Figures, the Circle and Proportion. By J. M. WILSON, M.A., Fellow of St. John's College, Cambridge, and Mathematical Master in Rugby School. Extra fcap. 8vo. 3*s.* 6*d.*

"*It is an actual substitute for the first two books of Euclid, in which many of his propositions are drawn out from the conception of straightness, parallelism, angles, with wonderful ease and simplicity, and the methods employed have the great merit of suggesting a ready application to the solution of fresh problems.*"—GUARDIAN.

ELEMENTARY GEOMETRY. PART II. (separately). The Circle and Proportion. By J. M. WILSON, M.A. Extra fcap. 8vo. 2*s.* 6*d.*

Wilson (W. P.) — A TREATISE ON DYNAMICS. By W. P. WILSON, M.A., Fellow of St. John's College, Cambridge, and Professor of Mathematics in Queen's College, Belfast. 8vo. 9*s.* 6*d.*

Wolstenholme. — A BOOK OF MATHEMATICAL PROBLEMS, on Subjects included in the Cambridge Course. By JOSEPH WOLSTENHOLME, Fellow of Christ's College, sometime Fellow of St. John's College, and lately Lecturer in Mathematics at Christ's College. Crown 8vo. cloth. 8*s.* 6*d.*

CONTENTS:—*Geometry (Euclid)—Algebra—Plane Trigonometry—Geometrical Conic Sections—Analytical Conic Sections—Theory of Equations—Differential Calculus—Integral Calculus—Solid Geometry—Statics—Elementary Dynamics—Newton—Dynamics of a Point—Dynamics of a Rigid Body—Hydrostatics—Geometrical Optics—Spherical Trigonometry and Plane Astronomy.*

SCIENCE.

The importance of Science as an element of sound education is now generally acknowledged; and accordingly it is obtaining a prominent place in the ordinary course of school instruction. It is the intention of the Publishers to produce a complete series of Scientific Manuals, affording full and accurate elementary information, conveyed in clear and lucid English. The authors are well known as among the foremost men of their several departments; and their names form a ready guarantee for the high character of the books. Subjoined is a list of those manuals that have already appeared, with a short account of each. Others are in active preparation; and the whole will constitute a standard series specially adapted to the requirements of beginners, whether for private study or for school instruction.

ASTRONOMY, by the Astronomer Royal.

POPULAR ASTRONOMY. With Illustrations. By G. B. AIRY, Astronomer Royal. Sixth and cheaper Edition. 18mo. cloth. 4*s*. 6*d*.

This work consists of six lectures, which are intended "to explain to intelligent persons the principles on which the instruments of an Observatory are constructed (omitting all details, so far as they are merely sub-

sidiary), and the principles on which the observations made with these instruments are treated for deduction of the distances and weights of the bodies of the Solar System, and of a few stars, omitting all minutiæ of formulæ, and all troublesome details of calculation." The speciality of this volume is the direct reference of every step to the Observatory, and the full description of the methods and instruments of observation.

ASTRONOMY.

MR. LOCKYER'S ELEMENTARY LESSONS IN ASTRONOMY. With Coloured Diagram of the Spectra of the Sun, Stars, and Nebulæ, and numerous Illustrations. By J. NORMAN LOCKYER, F.R.A.S. Third Thousand. 18mo. 5*s.* 6*d.*

The author has here aimed to give a connected view of the whole subject, and to supply facts, and ideas founded on the facts, to serve as a basis for subsequent study and discussion. The chapters treat of the Stars and Nebulæ; the Sun; the Solar System; Apparent Movements of the Heavenly Bodies; the Measurement of Time; Light; the Telescope and Spectroscope; Apparent Places of the Heavenly Bodies; the Real Distances and Dimensions; Universal Gravitation. The most recent astronomical discoveries are incorporated. Mr. Lockyer's work supplements that of the Astronomer Royal mentioned in the previous article.

PHYSIOLOGY.

PROFESSOR HUXLEY'S LESSONS IN ELEMENTARY PHYSIOLOGY. With numerous Illustrations. By T. H. HUXLEY, F.R.S. Professor of Natural History in the Royal School of Mines. Eighth Thousand. 18mo. cloth. 4*s.* 6*d.*

This book describes and explains, in a series of graduated lessons, the principles of Human Physiology; or the Structure and Functions of the Human Body. The first lesson supplies a general view of the subject. This is followed by sections on the Vascular or Veinous System, and the Circulation; the Blood and the Lymph; Respiration; Sources of Loss and of Gain to the Blood; the Function of Alimentation; Motion and Locomotion; Sensations and Sensory Organs; the Organ of Sight; the Coalescence of Sensations with one another and with other States of Consciousness; the Nervous System and Innervation; Histology, or the Minute Structure of the Tissues. A Table of Anatomical and Physiological Constants is appended. The lessons are fully illustrated by numerous engravings. The manual is primarily intended to serve as a text-book for teachers and learners in boys' and girls' schools.

QUESTIONS ON HUXLEY'S PHYSIOLOGY FOR SCHOOLS. By T. ALCOCK, M.D. 18mo. 1*s.* 6*d.*

These Questions were drawn up as aids to the instruction of a class of young people in Physiology.

BOTANY.

PROFESSOR OLIVER'S LESSONS IN ELEMENTARY BOTANY. With nearly Two Hundred Illustrations. By DANIEL OLIVER, F.R.S., F.L.S. Sixth Thousand. 18mo. cloth, 4*s.* 6*d.*

This book is designed to teach the Elements of Botany on Professor Henslow's plan of selected Types and by the use of Schedules. The earlier chapters, embracing the elements of Structural and Physiological Botany, introduce us to the methodical study of the Ordinal Types. The concluding chapters are entitled, "How to dry Plants" and "How to describe Plants." A valuable Glossary is appended to the volume. In the preparation of this work free use has been made of the manuscript materials of the late Professor Henslow.

CHEMISTRY.

PROFESSOR ROSCOE'S LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC. By HENRY E. ROSCOE, F.R.S., Professor of Chemistry in Owens College, Manchester. With numerous Illustrations and Chromo-Litho. of the Solar Spectrum, and of the Alkalies and Alkaline Earths, *New Edition.* Thirteenth Thousand. 18mo. cloth. 4*s.* 6*d.*

It has been the endeavour of the author to arrange the most important facts and principles of Modern Chemistry in a plain but concise and scientific form, suited to the present requirements of elementary instruction. For the purpose of facilitating the attainment of exactitude in the knowledge of the subject, a series of exercises and questions upon the lessons have been added. The metric system of weights and measures, and the centigrade thermometric scale, are used throughout the work. The new Edition, besides new wood-cuts, contains many additions and improvements, and includes the most important of the latest discoveries.

MISCELLANEOUS.

ATLAS OF EUROPE. GLOBE EDITION. Uniform in size with Macmillan's Globe Series, containing 45 Coloured Maps, on a uniform scale and projection: with Plans of London and Paris, and a copious Index. Strongly bound in half-morocco, with flexible back, 9*s.*

This Atlas includes all the countries of Europe in a series of 48 Maps drawn on the same scale, with an Alphabetical Index to the situation of more than ten thousand places; and the relation of the various maps and countries to each other is defined in a general Key-map. The identity of scale in all the maps facilitates the comparison of extent and distance, and conveys a just impression of the magnitude of different countries. The size suffices to show the provincial divisions, the railways and main roads, the principal rivers and mountain ranges. "This atlas," writes the British Quarterly, *"will be an invaluable boon for the school, the desk, or the traveller's portmanteau."*

Bates & Lockyer.—A CLASS-BOOK OF GEOGRAPHY. Adapted to the recent Programme of the Royal Geographical Society. By H. W. BATES, Assistant Secretary to the Royal Geographical Society, and J. N. LOCKYER, F.R.A.S.

[*In the Press.*

CAMEOS FROM ENGLISH HISTORY. From Rollo to Edward II. By the Author of "The Heir of Redclyffe." Extra fcap. 8vo. Second Edition. 5*s.*

The endeavour has not been to chronicle facts, but to put together a series of pictures of persons and events, so as to arrest the attention, and give some individuality and distinctness to the recollection, by gathering together details at the most memorable moments. The "Cameos" are intended as a book for young people just beyond the elementary histories of England,

and able to enter in some degree into the real spirit of events, and to be struck with characters and scenes presented in some relief. "Instead of dry details," says the Nonconformist, *"we have living pictures, faithful, vivid, and striking."*

HISTORICAL SELECTIONS. Readings from the best Authorities on English and European History. Selected and Arranged by E. M. SEWELL and C. M. YONGE. Crown 8vo. 6*s*.

When young children have acquired the outlines of History from abridgments and catechisms, and it becomes desirable to give a more enlarged view of the subject, in order to render it really useful and interesting, a difficulty often arises as to the choice of books. Two courses are open, either to take a general and consequently dry history of facts, such as Russel's Modern Europe, or to choose some work treating of a particular period or subject, such as the works of Macaulay and Froude. The former course usually renders history uninteresting; the latter is unsatisfactory, because it is not sufficiently comprehensive. To remedy this difficulty selections, continuous and chronological, have, in the present volume, been taken from the larger works of Freeman, Milman, Palgrave, and others, which may serve as distinct landmarks of historical reading. "We know of scarcely anything," says the Guardian, *of this volume, "which is so likely to raise to a higher level the average standard of English education."*

Hole.—A GENEALOGICAL STEMMA OF THE KINGS OF ENGLAND AND FRANCE. By the Rev. C. HOLE. On Sheet. 1*s*.

The different families are printed in distinguishing colours, thus facilitating reference.

A BRIEF BIOGRAPHICAL DICTIONARY. Compiled and Arranged by CHARLES HOLE, M.A., Trinity College, Cambridge. Second Edition, 18mo. neatly and strongly bound in cloth. 4*s*. 6*d*.

The most comprehensive Biographical Dictionary in the world, containing more than 18,000 *names of persons of all countries, with dates of birth and death, and what they were distinguished for. Reference is also made to Biographies of the persons mentioned. Extreme care has been bestowed on the verification of the dates, and thus numerous errors, current in previous works, have been corrected.*

"An invaluable addition to our manuals of reference, and, from its moderate price, cannot fail to become as popular as it is useful."—TIMES.

Jephson.—SHAKESPEARE'S TEMPEST. With Glossarial and Explanatory Notes. By the Rev. J. M. JEPHSON. 18mo. 1*s*. 6*d*.

It is important to find some substitute for classical study, and it is believed that such a substitute may be found in the Plays of Shakespeare. Each sentence of Shakespeare becomes, like a sentence in Thucydides or Cicero, a lesson in the origin and derivation of words, and in the fundamental rules of grammatical construction. On this principle the present edition of the "Tempest" has been prepared. The text is taken from the "Cambridge Shakespeare."

Oppen.—FRENCH READER. For the Use of Colleges and Schools. Containing a graduated Selection from modern Authors in Prose and Verse; and copious Notes, chiefly Etymological. By EDWARD A. OPPEN. Fcap. 8vo. cloth. 4*s*. 6*d*.

This is a Selection from the best modern authors of France. Its distinctive feature consists in its etymological notes, connecting French with the classical and modern languages, including the Celtic. This subject has hitherto been little discussed even by the best-educated teachers.

A SHILLING BOOK OF GOLDEN DEEDS. A Reading Book for Schools and General Readers. By the Author of "The Heir of Redclyffe." 18mo. cloth.

A record of some of the good and great deeds of all time, abridged from the larger work of the same author in the Golden Treasury Series.

Vaughan, C. M. — A SHILLING BOOK OF WORDS FROM THE POETS. By C. M. VAUGHAN. 18mo. cloth.

It has been felt of late years that the children of our parochial schools, and those classes of our countrymen which they commonly represent, are capable of being interested, and therefore benefited also, by something higher in the scale of poetical composition than those brief and somewhat puerile fragments to which their knowledge was formerly restricted. An attempt has here been made to supply the want by forming a selection at once various and unambitious; healthy in tone, just in sentiment, elevating in thought, and beautiful in expression.

Thring.—Works by EDWARD THRING, M.A., Head Master of Uppingham.

THE ELEMENTS OF GRAMMAR TAUGHT IN ENGLISH, with Questions. Fourth Edition. 18mo. 2*s.*

This little work is chiefly intended for teachers and learners. It took its rise from questionings in National Schools, and the whole of the first part is merely the writing out in order the answers to questions which have been used already with success. A chapter on Learning Language is especially addressed to teachers.

THE CHILD'S GRAMMAR. Being the Substance of "The Elements of Grammar taught in English," adapted for the Use of Junior Classes. A New Edition. 18mo. 1*s.*

SCHOOL SONGS. A Collection of Songs for Schools. With the Music arranged for four Voices. Edited by the Rev. E. THRING and H. RICCIUS. Folio. 7*s.* 6*d.*

There is a tendency in schools to stereotype the forms of life. Any genial solvent is valuable. Games do much; but games do not penetrate to domestic life, and are much limited by age. Music supplies the want. The collection includes the "Agnus Dei," Tennyson's "Light Brigade," Macaulay's "Ivry," &c. among other pieces.

Trench, Archbishop.—HOUSEHOLD BOOK OF ENGLISH POETRY. Selected and Arranged, with Notes, by R. C. TRENCH, D.D., Archbishop of Dublin. Extra fcap. 8vo. 5*s.* 6*d.*

This volume is called a "Household Book," by this name implying that it is a book for all—that there is nothing in it to prevent it from being confidently placed in the hands of every member of the household. Specimens of all classes of poetry are given, including selections from living authors. The Editor has aimed to produce a book "which the emigrant, finding room for little not absolutely necessary, might yet find room for it in his trunk, and the traveller in his knapsack, and that on some narrow shelves where there are few books this might be one."

"The Archbishop has conferred in this delightful volume an important gift on the whole English-speaking population of the world."—PALL MALL GAZETTE.

DIVINITY

Wright, W. Aldis (Co-Editor of the "Cambridge Shakespeare"). —THE BIBLE WORD-BOOK. A Glossary of Old English Bible Words. By J. EASTWOOD, M.A., of St. John's College, and W. ALDIS WRIGHT, M.A., Trinity College, Cambridge. 18mo. 5*s.* 6*d.*

It is the object of this Glossary to explain and illustrate all such words, phrases, and constructions, in the Authorized Version of the Old and New Testaments and the Apocrypha, and in the Book of Common Prayer, as are either obsolete or archaic. Full explanations are supplied, and these illustrated by numerous citations from the elder writers. An index of editions quoted is appended. Apart from its immediate subject, this work serves to illustrate a well-marked period in the history of the English language. It is thus of distinct philological value.

Hardwick.—A HISTORY OF THE CHRISTIAN CHURCH. Middle Age. From Gregory the Great to the Excommunication of Luther. By ARCHDEACON HARDWICK. Edited by FRANCIS PROCTER, M.A. With Four Maps constructed for this work by A. KEITH JOHNSTON. Second Edition. Crown 8vo. 10*s.* 6*d.*

The ground-plan of this treatise coincides in many points with one adopted at the close of the last century in the colossal work of Schröckh, and since that time by others of his thoughtful countrymen; but in arranging the materials a very different course has frequently been pursued. With regard to the opinions of the author, he is willing to avow distinctly that he has construed history with the specific prepossessions of an Englishman and a member of the English Church. The reader is constantly referred to the authorities, both original and critical, on which the statements are founded.

Hardwick, Archdeacon—*continued.*

A HISTORY OF THE CHRISTIAN CHURCH DURING THE REFORMATION. By ARCHDEACON HARDWICK. Revised by FRANCIS PROCTER, M.A. Second Edition. Crown 8vo. 10*s*. 6*d*.

This volume is intended as a sequel and companion to the "History of the Christian Church during the Middle Age." The author's earnest wish has been to give the reader a trustworthy version of those stirring incidents which mark the Reformation period, without relinquishing his former claim to characterise peculiar systems, persons, and events according to the shades and colours they assume, when contemplated from an English point of view, and by a member of the Church of England.

Maclear.—Works by the Rev. G. F. MACLEAR, B.D., Head Master of King's College School, and Preacher at the Temple Church.

A CLASS-BOOK OF OLD TESTAMENT HISTORY. Fourth Edition, with Four Maps. 18mo. cloth. 4*s*. 6*d*.

This volume forms a Class-book of Old Testament History from the earliest times to those of Ezra and Nehemiah. In its preparation the most recent authorities have been consulted, and wherever it has appeared useful, Notes have been subjoined illustrative of the Text, and, for the sake of more advanced students, references added to larger works. The Index has been so arranged as to form a concise dictionary of the persons and places mentioned in the course of the narrative; while the maps, which have been prepared with considerable care at Stanford's Geographical Establishment, will, it is hoped, materially add to the value and usefulness of the Book.

A CLASS-BOOK OF NEW TESTAMENT HISTORY, including the Connexion of the Old and New Testament. With Four Maps. Third Edition. 18mo. cloth. 5*s*. 6*d*.

A sequel to the author's Class-book of Old Testament History, continuing the narrative from the point at which it there ends, and carrying it on to the close of St. Paul's second imprisonment at Rome. In its preparation, as in that of the former volume, the most recent and trustworthy authorities have been consulted, notes subjoined, and references to larger works added. It is thus hoped that it may prove at once an useful class-book and a convenient companion to the study of the Greek Testament.

Maclear, Rev. G. F.—*continued.*

A SHILLING BOOK OF OLD TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo. cloth.

A SHILLING BOOK OF NEW TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo. cloth.

These works have been carefully abridged from the author's larger manuals.

CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. Second Edition. 18mo. cloth. 2*s.* 6*d.*

This may be regarded as a sequel to the Class-books of Old and New Testament History. Like them, it is furnished with notes and references to larger works, and it is hoped that it may be found, especially in the higher forms of our Public Schools, to supply a suitable manual of Instruction in the chief doctrines of the English Church, and a useful help in the preparation of Candidates for Confirmation.

A FIRST CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND, with Scripture Proofs, for Junior Classes and Schools. 18mo. 6*d.*

THE ORDER OF CONFIRMATION. A Sequel to the Class Book of the Catechism. For the use of Candidates for Confirmation. With Prayers and Collects. 18mo. 3*d.*

Procter.—A HISTORY OF THE BOOK OF COMMON PRAYER, with a Rationale of its Offices. By FRANCIS PROCTER, M.A. Eighth Edition, revised and enlarged. Crown 8vo. 10*s.* 6*d.*

In the course of the last twenty years the whole question of Liturgical knowledge has been reopened with great learning and accurate research, and it is mainly with the view of epitomizing extensive publications, and correcting the errors and misconceptions which had obtained currency, that the present volume has been put together.

Procter and Maclear.—AN ELEMENTARY INTRODUCTION TO THE BOOK OF COMMON PRAYER. Third Edition, re-arranged and supplemented by an Explanation of the Morning and Evening Prayer and the Litany. By the Rev. F. PROCTER and the Rev. G. F. MACLEAR. Third Edition. 18mo. 2*s*. 6*d*.

As in the other Class-books of the series, notes have also been subjoined, and references given to larger works, and it is hoped that the volume will be found adapted for use in the higher forms of our Public Schools, and a suitable manual for those preparing for the Oxford and Cambridge local examinations. This new Edition has been considerably altered, and several important additions have been made. Besides a re-arrangement of the work generally, the Historical Portion has been supplemented by an Explanation of the Morning and Evening Prayer and of the Litany.

PSALMS OF DAVID CHRONOLOGICALLY ARRANGED. BY FOUR FRIENDS. An Amended Version, with Historical Introduction and Explanatory Notes. Crown 8vo. 10*s*. 6*d*.

To restore the Psalter as far as possible to the order in which the Psalms were written,—to give the division of each Psalm into strophes, of each strophe into the lines which composed it,—to amend the errors of translation, is the object of the present Edition. Professor Ewald's works, especially that on the Psalms, have been extensively consulted.

This book has been used with satisfaction by masters for private work in higher classes in schools.

Ramsay.—THE CATECHISER'S MANUAL; or, the Church Catechism illustrated and explained, for the use of Clergymen, Schoolmasters, and Teachers. By the Rev. ARTHUR RAMSAY, M.A. Second Edition. 18mo. 1*s*. 6*d*.

A clear explanation of the Catechism, by way of question and answer.

Simpson.—AN EPITOME OF THE HISTORY OF THE CHRISTIAN CHURCH. By WILLIAM SIMPSON, M.A. Fourth Edition. Fcap. 8vo. 3*s*. 6*d*.

A compendious summary of Church History.

Swainson.—A HANDBOOK to BUTLER'S ANALOGY. By C. A. SWAINSON, D.D., Norrisian Professor of Divinity at Cambridge. Crown 8vo. 1*s.* 6*d.*

This manual is designed to serve as a handbook or road-book to the Student in reading the Analogy, to give the Student a sketch or outline map of the country on which he is entering, and to point out to him matters of interest as he passes along.

Westcott.—A GENERAL SURVEY OF THE HISTORY OF THE CANON OF THE NEW TESTAMENT DURING THE FIRST FOUR CENTURIES. By BROOKE FOSS WESTCOTT, B.D., Canon of Peterborough. Second Edition, revised. Crown 8vo. 10*s.* 6*d.*

The Author has endeavoured to connect the history of the New Testament Canon with the growth and consolidation of the Church, and to point out the relation existing between the amount of evidence for the authenticity of its component parts, and the whole mass of Christian literature. Such a method of inquiry will convey both the truest notion of the connexion of the written Word with the living Body of Christ, and the surest conviction of its divine authority.

Of this work the Saturday Review *writes: "Theological students, and not they only, but the general public, owe a deep debt of gratitude to Mr. Westcott for bringing this subject fairly before them in this candid and comprehensive essay..... As a theological work it is at once perfectly fair and impartial, and imbued with a thoroughly religious spirit; and as a manual it exhibits, in a lucid form and in a narrow compass, the results of extensive research and accurate thought. We cordially recommend it."*

INTRODUCTION TO THE STUDY OF THE FOUR GOSPELS. By BROOKE FOSS WESTCOTT, B.D. Third Edition. Crown 8vo. 10*s.* 6*d.*

This book is intended to be an Introduction to the Study *of the Gospels. The author has made it a point carefully to study the researches of the great writers, and consciously to neglect none. There is an elaborate discussion appended "On the Primitive Doctrine of Inspiration."*

Westcott, Canon—*continued.*

A GENERAL VIEW OF THE HISTORY OF THE ENGLISH BIBLE. By BROOKE FOSS WESTCOTT, B.D. Crown 8vo. 10*s.* 6*d.*

"*The first trustworthy account we have had of that unique and marvellous monument of the piety of our ancestors.*"—DAILY NEWS.

THE BIBLE IN THE CHURCH. A Popular Account of the Collection and Reception of the Holy Scriptures in the Christian Churches. Second Edition. By BROOKE FOSS WESTCOTT, B.D. 18mo. cloth, 4*s.* 6*d.*

The present book is an attempt to answer a request, which has been made from time to time, to place in a simple form, for the use of general readers, the substance of the author's "History of the Canon of the New Testament." An elaborate and comprehensive Introduction is followed by chapters on the Bible of the Apostolic Age; on the Growth of the New Testament; the Apostolic Fathers; the Age of the Apologists: the First Christian Bible; the Bible Proscribed and Restored; the Age of Jerome and Augustine: the Bible of the Middle Ages in the West and in the East, and in the Sixteenth Century. Two appendices on the History of the Old Testament Canon before the Christian Era, and on the Contents of the most ancient MSS. of the Christian Bible, complete the volume.

THE GOSPEL OF THE RESURRECTION. Thoughts on its Relation to Reason and History. By BROOKE FOSS WESTCOTT, B.D. New Edition. Fcap. 8vo. 4*s.* 6*d.*

This Essay is an endeavour to consider some of the elementary truths of Christianity as a miraculous Revelation from the side of History and Reason. If the arguments which are here adduced are valid, they will go far to prove that the Resurrection, with all that it includes, is the key to the history of man, and the complement of reason.

Wilson.—AN ENGLISH, HEBREW, AND CHALDEE LEXICON AND CONCORDANCE, to the more Correct Understanding of the English translation of the Old Testament, by reference to the Original Hebrew. By WILLIAM WILSON, D.D., Canon of Winchester, late Fellow of Queen's College, Oxford. Second Edition, carefully Revised. 4to. cloth. 25*s.*

The aim of this work is, that it should be useful to clergymen and all persons engaged in the study of the Bible, even when they do not possess a knowledge of Hebrew; while able Hebrew scholars have borne testimony to the help that they themselves have found in it.

BOOKS ON EDUCATION.

Arnold.—A FRENCH ETON; OR, MIDDLE CLASS EDUCATION AND THE STATE. By MATTHEW ARNOLD. Fcap. 8vo. cloth. 2*s*. 6*d*.

"*A very interesting dissertation on the system of secondary instruction in France, and on the advisability of copying the system in England.*"—SATURDAY REVIEW.

SCHOOLS AND UNIVERSITIES ON THE CONTINENT. 8vo. 10*s*. 6*d*.

The Author was in 1865 *charged by the Schools Inquiry Commissioners with the task of investigating the system of education for the middle and upper classes in France, Italy, Germany, and Switzerland. In the discharge of this task he was on the continent nearly seven months, and during that time he visited the four countries named, and made a careful study of the matters to which the Commissioners had directed his attention. The present volume contains the report which he made to them. It is here adapted to the use of the general reader.*

ESSAYS ON A LIBERAL EDUCATION. Edited by the Rev. F. W. FARRAR, M.A., F.R.S., Assistant Master at Harrow, late Fellow of Trinity College, Cambridge, and Hon. Fellow of King's College, London. Second Edition. 8vo. cloth. 10*s*. 6*d*.

CONTENTS:—*History of Classical Education, by Charles S. Parker, M.A.; Theory of Classical Education, by Henry Sedgwick, M.A.;*

Liberal Education in Universities, by John Seeley, M.A.; Teaching by means of Grammar, by E. E. Bowen, M.A.; Greek and Latin Verse-Composition, by the Rev. F. W. Farrar; Natural Science in Schools, by J. M. Wilson, M.A., F.G.S.; The Teaching of English, by J. W. Hales, M.A.; Education of the Reasoning Faculties, by W. Johnson, M.A.; The present Social Results of Classical Education, by Lord Houghton.

The Authors have sought to hasten the expansion and improvement of liberal education by showing in what light some of the most interesting questions of Educational Reform are viewed by men who have had opportunities for forming a judgment respecting them, and several of whom have been for some time engaged in the work of education at our Universities and Schools.

Farrar.—ON SOME DEFECTS IN PUBLIC SCHOOL EDUCATION. A Lecture delivered at the Royal Institution. With Notes and Appendices. Crown 8vo. 1*s.*

Jex-Blake.—A VISIT TO SOME AMERICAN SCHOOLS AND COLLEGES. By SOPHIA JEX-BLAKE. Crown 8vo. cloth. 6*s.*

"*In the following pages I have endeavoured to give a simple and accurate account of what I saw during a series of visits to some of the Schools and Colleges in the United States. . . . I wish simply to give other teachers an opportunity of seeing through my eyes what they cannot perhaps see for themselves, and to this end I have recorded just such particulars as I should myself care to know.*"—AUTHOR'S PREFACE.

"*Miss Blake gives a living picture of the Schools and Colleges themselves, in which that education is carried on.*"—PALL MALL GAZETTE.

Thring.—EDUCATION AND SCHOOL. By the Rev. EDWARD THRING, M.A., Head Master of Uppingham. Second Edition. Crown 8vo. cloth. 5*s.* 6*d.*

Youmans.—MODERN CULTURE: its True Aims and Requirements. A Series of Addresses and Arguments on the Claims of Scientific Education. Edited by EDWARD L. YOUMANS, M.D. Crown 8vo. 8*s.* 6*d.*

CONTENTS:—*Professor Tyndall on the Study of Physics; Dr. Daubeny on the Study of Chemistry; Professor Henfrey on the Study of Botany; Professor Huxley on the Study of Zoology; Dr. J. Paget on the Study of Physiology; Dr. Whewell on the Educational History of Science; Dr. Faraday on the Education of the Judgment; Dr. Hodgson on the Study of Economic Science; Mr. Herbert Spencer on Political Education; Professor Masson on College Education and Self Education; Dr. Youmans on the Scientific Study of Human Nature. An Appendix contains extracts from distinguished authors, and from the Scientific Evidence given before the Public Schools Commission.*

LONDON: R. CLAY, SONS, AND TAYLOR, PRINTERS.